

TITS ALTERNATIVE FOR CLOSED REAL ANALYTIC 4-MANIFOLDS OF NONPOSITIVE CURVATURE

XIANGDONG XIE

ABSTRACT. We study subgroups of fundamental groups of real analytic closed 4-manifolds with nonpositive sectional curvature. In particular, we are interested in the following question: if a subgroup of the fundamental group is not virtually free abelian, does it contain a free group of rank two? The technique involves the theory of general metric spaces of nonpositive curvature.

1. INTRODUCTION

In this paper we study subgroups of fundamental groups of real analytic closed 4-manifolds with nonpositive sectional curvature. One would like to know whether a subgroup contains a finite index free abelian subgroup if it does not contain a free group of rank two. The same question can be asked in the general setting of groups acting on Hadamard spaces. We recall a Hadamard space is a complete simply connected metric space with nonpositive curvature in the sense of A. D. Alexandrov. In general there is the following question (see Section 2.4 for the definition of a proper group action):

Tits Alternative. Let G be a group acting properly and cocompactly by isometries on a Hadamard space X . Is it true that every subgroup $H \subset G$ contains either a finite index free abelian subgroup or a free group of rank two?

The Tits alternative question has been answered affirmatively for the following spaces: trees ([PV]) or more generally Gromov hyperbolic Hadamard spaces ([G]); certain cubical complexes ([BSw]); Euclidean buildings of rank ≥ 3 or symmetric spaces ([T]); spaces with isolated flats ([HR]); certain square complexes ([X]). The Tits alternative question in the general case appears hard. In particular it is still open for Hadamard 4-manifolds and $CAT(0)$ 2-complexes. It is not even known ([Sw]) whether G has an infinite subgroup where each element is of finite order. S. Adams and W. Ballmann ([AB]) showed any amenable subgroup of the group G contains a finite index free abelian subgroup. When X is a piecewise smooth 2-complex where each edge is contained in at least two 2-cells, W. Ballmann and M. Brin ([BBr]) showed either G contains a free group of rank two or X is isometric to the Euclidean plane.

Recall a Hadamard manifold X has higher rank if each geodesic is contained in a 2-flat, that is, a convex subset of X isometric to $\mathbb{R}^2$; X has rank one otherwise. When the Hadamard space X is a Hadamard manifold and the group G is torsion free, the Tits alternative can be reduced to the case when X is an irreducible Hadamard manifold, thanks to Eberlein's results ([E1], [E2]) on lattices of reducible Hadamard manifolds. When X is an irreducible Hadamard manifold of higher rank, X is a higher rank symmetric space by the rank rigidity theorem ([B]) and the Tits alternative follows from Tits' theorem. Thus one only needs to consider rank one Hadamard manifolds.

The Tits alternative holds trivially for surfaces. It is also not hard to establish it for 3-manifolds, by using some nontrivial results on 3-manifold topology:

2000 *Mathematics Subject Classification.* Primary 53C20, 20F67; Secondary 20F65.

Key words and phrases. Tits alternative, rank one isometry, Hadamard space, nonpositive curvature, Tits metric.

Theorem 3.7. *The Tits alternative holds for subgroups of $\pi_1(M)$, where M is a closed 3-manifold with nonpositive sectional curvature.*

In this paper we mainly consider the case of real analytic closed 4-manifolds with nonpositive sectional curvature. Such manifolds have been studied by V. Schroeder ([S1]) and C. Hummel and V. Schroeder ([HS1], [HS2]). A higher rank submanifold in a Hadamard manifold X is a totally geodesic submanifold which is of higher rank as a Hadamard manifold. When $M = X/\Gamma$ is a real analytic closed 4-manifold with nonpositive sectional curvature, one possible class of higher rank submanifolds have the form $W = Q \times R$, where Q is a nonflat 2-dimensional Hadamard manifold. A *cycle* is a finite sequence $W_1, \dots, W_k$ of distinct higher rank submanifolds of the above form such that $W_i \cap W_{i+1} \neq \emptyset$ ($i = 1, 2, \dots, k-1$) and $W_k \cap W_1 \neq \emptyset$. Below is one of the main results of the paper.

Theorem 4.1. *The Tits alternative holds for subgroups of $\pi_1(M)$ if $M = X/\Gamma$ is a real analytic closed 4-manifold with nonpositive sectional curvature and there is no cycle in X .*

There exist real analytic closed 4-manifolds (see [AS]) satisfying the assumptions of the theorem.

Let $M = X/\Gamma$ be a real analytic closed 4-manifold with nonpositive sectional curvature. A *singular geodesic* in X is a geodesic c of the form $c = \{q\} \times R \subset Q \times R$, where $Q \times R$ is a higher rank submanifold and Q is a nonflat Hadamard 2-manifold. When two higher rank submanifolds $W_1 = Q_1 \times R$, $W_2 = Q_2 \times R$ intersect, the intersection is a 2-flat of the form $F = W_1 \cap W_2 = c_1 \times R$ where $c_1 \subset Q_1$ is a geodesic in Q_1 . The R directions in W_1 and W_2 give rise to two parallel families of singular geodesics in the 2 flat F . The angle between the two family is a *singular angle*. There are only a finite number of singular angles (see Section 2.3). U. Abresch and V. Schroeder ([AS]) have constructed a class of real analytic closed 4-manifolds with nonpositive sectional curvature where the singular angles are all equal to $\pi/2$.

Theorem 5.1. *The Tits alternative holds for subgroups of $\pi_1(M)$ if $M = X/\Gamma$ is a real analytic closed 4-manifold with nonpositive sectional curvature and all singular angles are equal to $\alpha > \frac{2}{5}\pi$.*

Although we are mainly interested in Hadamard manifolds in this paper, our proof uses general metric spaces with nonpositive curvature in the sense of A. D. Alexandrov. The proof of Theorem 4.1 is inspired by the JSJ decomposition in 3-manifold theory. Recall a Haken 3-manifold admits a JSJ decomposition and this decomposition induces a graph of groups decomposition for the fundamental group. It follows that the fundamental group acts on the Bass-Serre tree associated to the graph of groups. Notice that the Bass-Serre tree is not a Hadamard manifold and in general is not locally compact. In a similar way we decompose a real analytic closed 4-manifold M with nonpositive sectional curvature and construct a 2-complex associated to the decomposition. The fundamental group of M acts on the 2-complex by isometries. The 2-complex is a $CAT(-1)$ space (see Section 2.1 for definition) and should be considered as an analogue of the Bass-Serre tree.

The existence of free subgroups is closely related to the existence of rank one isometries. A rank one isometry is a hyperbolic isometry (see Section 2.2 for definition) g of a Hadamard space X such that no axis of g bounds a flat half plane, where a flat half plane is a convex subset of X isometric to the upper half plane: $\{(x, y) \in \mathbb{R}^2 : y \geq 0\}$. Each isometry of X induces a homeomorphism of the geometric boundary of X . If X is a $CAT(-1)$ space or a locally compact Hadamard space, then each rank one isometry of X acts on the geometric boundary in the same way as a hyperbolic isometry of the real hyperbolic space, see Theorem 2.11 or [B] and [G]. It follows that any group generated by two rank one isometries that do not share any fixed point in the geometric boundary contains a free group of rank two. Notice that it is not necessary to assume the action is proper.

For a general hyperbolic isometry, the dynamics of the induced homeomorphism on the geometric boundary have been studied by V. Schroeder ([BGS]) in the Hadamard manifold case and by K. Ruane ([R]) for locally compact Hadamard spaces. Let g be a hyperbolic isometry of a locally

compact Hadamard space X and $P(g)$ the parallel set of an axis of g , that is, $P(g)$ is the union of all geodesics that are parallel to an axis of g . Denote by $\partial_\infty X$ the geometric boundary of X . $P(g)$ is closed and convex in X and $\partial_\infty P(g)$ naturally embeds into $\partial_\infty X$. If $c : \mathbb{R} \rightarrow X$ is an axis of g , denote by $g(+\infty)$ and $g(-\infty)$ the points in $\partial_\infty X$ determined by the rays $c|_{[0,\infty)}$ and $c|_{(-\infty,0]}$ respectively. V. Schroeder and K. Ruane showed that for any $\xi \in \partial_\infty X - \partial_\infty P(g)$, the accumulation points of the orbit $\{g^i(\xi) : i \in \mathbb{Z}\}$ lie in $\partial_\infty P(g)$. K. Ruane ([R]) further proved the following result: if the Tits distance from $\xi \in \partial_\infty X$ to $g(-\infty)$ is greater than π , then there is a neighborhood U of ξ so that U is attracted to $g(+\infty)$ under the iteration of g . Here we still use g to denote the homeomorphism of the geometric boundary induced by g . Still the following question remains:

Question 1. *With the above notation. Is the following statement true: for any compact subset $K \subset \partial_\infty X - \partial_\infty P(g)$, and any neighborhood V of $\partial_\infty P(g)$, there is a positive integer N such that $g^n(K) \subset V$ for all $n \geq N$.*

If the answer to Question 1 is yes, then any group generated by two hyperbolic isometries f, g contains a free group of rank two whenever $\partial_\infty P(f)$ and $\partial_\infty P(g)$ (considered as subsets of $\partial_\infty X$) have empty intersection.

The existence of rank one isometries in a group is closely related to the dynamics of the group action on the limit set. While a lot is known about the group action on the limit set of discrete isometry groups of the real hyperbolic space, little is known for groups acting on Hadamard spaces or even Hadamard manifolds. Let G be a group acting properly on a Hadamard space such that the limit set (see Section 2.5 for definition) of G contains more than two points. The limit set is closed and G -invariant. If X is the real hyperbolic space, then the limit set is the only nonempty closed and G -invariant subset of the limit set. This is no longer the case in general, for instance, when $G = G_1 \times G_2$ and $X = X_1 \times X_2$ where $G_1 \subset \text{Isom}(X_1)$, $G_2 \subset \text{Isom}(X_2)$. It would be interesting to know when there are more than one closed and G -invariant subset. Recently W. Buyalo and W. Ballmann ([BB]) did some interesting work on the topic. In particular, using their arguments we can show the following:

Corollary 2.19. *Suppose Γ is a group of isometries of a locally compact Hadamard space X . If Γ does not contain any rank one isometry, then $\Lambda(\Gamma)$ has diameter at most 2π in the Tits metric.*

The paper is organized as follows. In Section 2 we recall basic facts about Hadamard spaces, and collect results that shall be needed later on. The topics covered in this section include: higher rank submanifolds in the universal covers of closed real analytic 4-manifolds with nonpositive sectional curvature and structure of Tits boundary of such manifolds, rank one isometry and free groups, action on the limit set. In Section 3 we establish the Tits alternative for closed 3-manifolds with nonpositive sectional curvature. In Section 4 we decompose closed real analytic 4-manifolds, construct the associated 2-complex and use the 2-complex to establish the main result of the paper (Theorem 4.1). In Section 5 we discuss the Tits alternative without assuming the nonexistence of cycles.

Acknowledgment. *I am grateful to Quo-Shin Chi for numerous discussions. I would also like to thank Bruce Kleiner, Blake Thornton, Kevin Scannell, Rachel Roberts, Victor Schroeder, Michael Kapovich and Alan Reid for remarks and communications on the subject.*

2. PRELIMINARIES

In this section we recall basic facts about Hadamard spaces, and collect results that shall be needed later on. We refer the reader to [B], [BGS], [BH], [S1] and [HS1] for more details on the material in this section.

2.1. $CAT(\kappa)$ Spaces. Although we are mainly interested in Hadamard manifolds in this paper, general metric spaces with upper curvature bounds will play an important role in the proof of our results. So here we recall the basic definitions below.

Let (X, d) be a metric space. A *geodesic* in X is a continuous map $\alpha : I \rightarrow X$ such that, for any point $t \in I$, there exists a neighborhood U of t with $d(\alpha(s_1), \alpha(s_2)) = |s_1 - s_2|$ for all $s_1, s_2 \in U$. If the above equality holds for all $s_1, s_2 \in I$, then we call α a *minimal geodesic*. The image of a geodesic shall also be called a geodesic. When I is a closed interval $[a, b]$, we say α is a geodesic segment of length $b - a$ and α connects $\alpha(a)$ and $\alpha(b)$. A metric space X is called a *geodesic metric space* if for any two points $x, y \in X$ there is a minimal geodesic segment connecting them.

A *triangle* in a metric space (X, d) is the union of three geodesic segments $\alpha_i : [a_i, b_i] \rightarrow X$ ($i = 1, 2, 3$) where $\alpha_1(b_1) = \alpha_2(a_2)$, $\alpha_2(b_2) = \alpha_3(a_3)$ and $\alpha_3(b_3) = \alpha_1(a_1)$. For any real number κ , let M_κ^2 stand for the 2-dimensional simply connected complete Riemannian manifold with constant curvature κ , and $D(\kappa)$ denote the diameter of M_κ^2 ($D(\kappa) = \infty$ if $\kappa \leq 0$). Given a triangle $\Delta = \alpha_1 \cup \alpha_2 \cup \alpha_3$ in a metric space X where $\alpha_i : [a_i, b_i] \rightarrow X$ ($i = 1, 2, 3$), a triangle Δ' in M_κ^2 is a *comparison triangle* for Δ if they have the same edge lengths, that is, if $\Delta' = \alpha'_1 \cup \alpha'_2 \cup \alpha'_3$ and $\alpha'_i : [a_i, b_i] \rightarrow M_\kappa^2$ ($i = 1, 2, 3$). A point $x' \in \Delta'$ corresponds to a point $x \in \Delta$ if there is some i and some $t_i \in [a_i, b_i]$ with $x' = \alpha'_i(t_i)$ and $x = \alpha_i(t_i)$. We notice if the perimeter of a triangle $\Delta = \alpha_1 \cup \alpha_2 \cup \alpha_3$ in X is less than $2D(\kappa)$, that is, if $length(\alpha_1) + length(\alpha_2) + length(\alpha_3) < 2D(\kappa)$, then there is a unique comparison triangle (up to isometry) in M_κ^2 for Δ .

Definition 2.1. A complete metric space X is called a $CAT(\kappa)$ space if

- (i) every two points $x_1, x_2 \in X$ with $d(x_1, x_2) < D(\kappa)$ are connected by a minimal geodesic segment;
- (ii) for any triangle Δ in X with perimeter less than $2D(\kappa)$ and any two points $x, y \in \Delta$, the inequality $d(x, y) \leq d(x', y')$ holds, where x' and y' are the points on a comparison triangle for Δ corresponding to x and y respectively.

A complete metric space has *curvature* $\leq \kappa$ if each point has a $CAT(\kappa)$ neighborhood.

A simply connected complete Riemannian manifold with sectional curvature $\leq \kappa$ ($\kappa \leq 0$) is $CAT(\kappa)$. Simplicial metric trees and more generally R -trees are $CAT(\kappa)$ for any κ . The following lemma follows from the above definition.

Lemma 2.2. Let X be a $CAT(1)$ space and $\alpha : [a, b] \rightarrow X$ a geodesic with $a < b$ and $\alpha(a) = \alpha(b)$. Then the length of α is at least 2π .

2.2. Hadamard Spaces and Their Ideal Boundaries. A $CAT(0)$ space is also called a *Hadamard space*. Let (X, d) be a Hadamard space. Then the distance function $d : X \times X \rightarrow \mathbb{R}$ is convex, there is a unique geodesic segment between any two points and X is contractible. For any $x, y \in X$, xy denotes the unique geodesic segment connecting x and y . A *ray* starting from $p \in X$ is a geodesic $c : [0, \infty) \rightarrow X$ with $c(0) = p$. Two rays c_1 and c_2 are *asymptotic* if $d(c_1(t), c_2(t))$ is a bounded function on the interval $[0, \infty)$. The *ideal boundary* of X is the set ∂X of asymptotic classes of rays in X . For any $p \in X$ and any $\xi \in \partial X$, there is a unique ray (denoted by $p\xi$) that starts from p and belongs to ξ . Thus for any $p \in X$ we can identify ∂X with the set of rays starting from p . Let c and c_i ($i = 1, 2, \dots$) be rays starting from p ; we say $\{c_i\}_{i=1}^\infty$ converges to c if c_i converges to c uniformly on compact subsets of $[0, \infty)$. Similarly for $x_i \in X$ ($i = 1, 2, \dots$), we say $\{x_i\}_{i=1}^\infty$ converges to $\xi \in X \cup \partial X$ if px_i converges to $p\xi$ uniformly on compact subsets. In this way we define a topology on $X \cup \partial X$. It is easy to check that this topology is independent of the point $p \in X$. Both this topology and the induced topology on ∂X are called the *cone topology*. The topology on X induced by the cone topology coincides with the metric topology on X . ∂X together with the cone topology is called the *geometric boundary* of X , and denoted by $\partial_\infty X$. We set $\overline{X} = X \cup \partial_\infty X$.

The Tits metric on the ideal boundary is defined as follows. Let $\xi, \eta \in \partial_\infty X$. For $p \in X$, let $c_1(t)$ and $c_2(t)$ be the rays that start from p and asymptotic to ξ and η respectively. The Tits angle

$\angle_T(\xi, \eta)$ between ξ and η is given by:

$$\sin\left(\frac{\angle_T(\xi, \eta)}{2}\right) = \lim_{t \rightarrow \infty} \frac{d(c_1(t), c_2(t))}{2t}.$$

This definition is independent of the point p . The *Tits metric* d_T is the path metric induced by $\angle_T$. In particular, if $\xi, \eta \in \partial_\infty X$ are in different Tits components, then $d_T(\xi, \eta) = \infty$. $\partial_\infty X$ with the Tits metric d_T is called the *Tits boundary* of X and denoted by $\partial_T X$. The Tits topology and the cone topology are generally quite different.

Below we collect some basic facts concerning the Tits metric. For more details please see [BGS] and [BH]. For any geodesic $c : R \rightarrow X$ in a Hadamard space, we call the two points in $\partial_\infty X$ determined by the two rays $c|_{[0, +\infty)}$ and $c|_{(-\infty, 0]}$ the endpoints of c , and denote them by $c(+\infty)$ and $c(-\infty)$ respectively.

Proposition 2.3. *Let X be a Hadamard space and $\xi, \eta \in \partial_T X$.*

- (i) $\partial_T X$ is a CAT(1) space;
- (ii) *If X is locally compact and $d_T(\xi, \eta) > \pi$, then there is a geodesic in X with ξ, η as endpoints;*
- (iii) *If X is locally compact and $d_T(\xi, \eta) < \infty$, then there is a minimal geodesic in $\partial_T X$ from ξ to η .*

Let X be a Hadamard space and $g : X \rightarrow X$ an isometry of X . g is called a *hyperbolic isometry* if it translates a geodesic, that is, if there is a geodesic $c : R \rightarrow X$ and a positive number l so that $g(c(t)) = c(t+l)$ for all $t \in R$; the geodesic c is called an *axis* of g . All the axes of g are parallel, thus it makes sense to denote $g(+\infty) = c(+\infty)$, $g(-\infty) = c(-\infty)$. Recall two geodesics $c_1, c_2 : R \rightarrow X$ are parallel if $d(c_1(t), c_2(t))$ is a bounded function over R .

For a hyperbolic isometry g , let $\text{Min}(g)$ be the union of all the axes of g . The subset $\text{Min}(g) \subset X$ is closed and convex in X and splits isometrically as $Y \times R$, where each $\{y\} \times R$ ($y \in Y$) is an axis of g . The geometric boundary $\partial_\infty \text{Min}(g)$ naturally embeds into $\partial_\infty X$.

Each isometry g of X induces a homeomorphism of the geometric boundary $\partial_\infty X$, which we still denote by g .

Theorem 2.4. ([R]) *Let X be a Hadamard space and g a hyperbolic isometry of X . Then the fixed point set of g on $\partial_\infty X$ is $\partial_\infty \text{Min}(g)$.*

2.3. Real Analytic 4-manifolds With Nonpositive Sectional Curvature. In this section we recall some facts concerning the universal covers of real analytic closed 4-manifolds with nonpositive sectional curvature. The reader is referred to [S1], [HS1] and [AS] for more details.

Let X be a Hadamard manifold, i.e., a simply connected complete Riemannian manifold with nonpositive sectional curvature. A k -flat in X is a totally geodesic submanifold isometric to the k -dimensional Euclidean space $\mathbb{E}^k$. We say X has *higher rank* if each geodesic in X is contained in a 2-flat, and X has *rank 1* otherwise. A complete totally geodesic submanifold of X is a *higher rank submanifold* if it has higher rank as a Hadamard manifold. A *maximal higher rank submanifold* of X is a higher rank submanifold that is maximal with respect to inclusion.

Let M be a closed Riemannian manifold with nonpositive sectional curvature and X its universal cover. Then $M = X/\Gamma$ where Γ is the group of deck transformations acting on X as isometries. A complete totally geodesic submanifold W of X is *closed* if $W/\text{Stab}_\Gamma W$ is compact, where $\text{Stab}_\Gamma W = \{\gamma \in \Gamma : \gamma(W) = W\}$ is the stabilizer of W in Γ . We say M is of rank 1 if X is of rank 1.

Theorem 2.5. ([S1]) *Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature. Then:*

- (i) *each maximal higher rank submanifold of X is isometric to one of the following: $\mathbb{E}^2$, $\mathbb{E}^3$, $Q \times R$, where Q is a nonflat 2-dimensional Hadamard manifold;*
- (ii) *each maximal higher rank submanifold is closed;*
- (iii) *there are only a finite number of maximal higher rank submanifolds modulo Γ ;*
- (iv) *$W_1 \cap W_2 \cap W_3 = \emptyset$ for any three distinct 3-dimensional higher rank submanifolds W_1, W_2, W_3 of X .*

For $M = X/\Gamma$ as in Theorem 2.5, we let $\mathcal{W}$ be the set of maximal higher rank submanifolds of X that are of the form $Q \times R$ for nonflat 2-dimensional Hadamard manifolds Q .

Theorem 2.6. ([S1]) *Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature, and W_1, W_2 maximal higher rank submanifolds of X with $W_1 \cap W_2 \neq \emptyset$. Then one of the following holds:*

- (i) W_1, W_2 are both isometric to $\mathbb{E}^2$. In this case $W_1 \cap W_2$ is a point;
- (ii) $W_1, W_2 \in \mathcal{W}$. In this case, W_1 and W_2 are perpendicular to each other, $W_1 \cap W_2$ is a 2-flat and $W_1 \cap W_2 = c_1 \times R \subset Q_1 \times R = W_1$ for a geodesic c_1 of Q_1 .

Notice for a complete totally geodesic submanifold W of X , the geometric boundary $\partial_\infty W$ naturally embeds into $\partial_\infty X$.

Theorem 2.7. ([HS1]) *Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature, and $\partial_T X$ the Tits boundary of X . Assume $\mathcal{C}$ is a connected component of $\partial_T X$. Then exactly one of the following statements is true:*

- (i) $\mathcal{C}$ consists of a single point;
- (ii) $\mathcal{C} = \partial_\infty F$, where F is a 2-flat;
- (iii) $\mathcal{C} = \partial_\infty F$, where F is a 3-flat;
- (iv) $\mathcal{C} = \bigcup_{W \in \mathcal{W}^*} \partial_\infty W$, where $\mathcal{W}^* \subset \mathcal{W}$ is a subset such that $\bigcup_{W \in \mathcal{W}^*} W$ is a connected component of $\bigcup_{W \in \mathcal{W}} W \subset X$;
- (v) $\mathcal{C}$ is isometric to a closed interval with length $< \pi$.

For convenience, we introduce the following definition.

Definition 2.8. Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature. An n -cycle in X is a sequence of n distinct higher rank submanifolds $W_1, W_2, \dots, W_n$ such that all $W_i \in \mathcal{W}$ and $W_i \cap W_{i+1} \neq \emptyset$ for all i , where indices are taken modulo n . A cycle in X is an n -cycle for some n .

Proposition 2.9. ([HS1]) *Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature. If there is an n -cycle in X , then $n \geq 5$.*

For any metric space Z and any $A \subset Z$, $\epsilon > 0$, the ϵ -neighborhood of A is $N_\epsilon(A) = \{z \in Z : d(z, a) < \epsilon \text{ for some } a \in A\}$. Let X be a Hadamard manifold. Two complete totally geodesic submanifolds H_1 and H_2 of X are parallel if there is some $\epsilon > 0$ with $H_2 \subset N_\epsilon(H_1)$ and $H_1 \subset N_\epsilon(H_2)$. Let H be a complete totally geodesic submanifold of X , the parallel set P_H of H is the union of all complete totally geodesic submanifolds of X that are parallel to H . P_H is a closed convex subset of X and splits isometrically as $P_H = H \times Y$, where each $H \times \{y\}$ ($y \in Y$) is a complete totally geodesic submanifold parallel to H . In general P_H has boundary and is not a manifold. When the Riemannian manifold X is real analytic, P_H is complete totally geodesic.

2.4. Rank One Isometries and Free Groups. Let X be a Hadamard space. A *flat half plane* in X is the image of an isometric embedding $f : \{(x, y) \in \mathbb{E}^2 : y \geq 0\} \rightarrow X$, and in this case we say the geodesic $c : R \rightarrow X$, $c(t) = f(t, 0)$ bounds the flat half plane.

Definition 2.10. A hyperbolic isometry g of a Hadamard space X is called a *rank one isometry* if no axis of g bounds a flat half plane.

If X is a Gromov hyperbolic Hadamard space, then each hyperbolic isometry of X is rank one. We note a $CAT(-1)$ space is Gromov hyperbolic. The following theorem is due to W. Ballmann in the case of locally compact Hadamard spaces and to M. Gromov in the case of Gromov hyperbolic spaces. Recall an isometry g of a Hadamard space X induces a homeomorphism of $\overline{X}$, which we still denote by g .

Theorem 2.11. ([B], [G]) *Let X be a Hadamard space that is either locally compact or Gromov hyperbolic, and g a rank one isometry of X . Given any neighborhoods U of $g(+\infty)$ and V of $g(-\infty)$ in $\overline{X}$, there is an $n \geq 0$ such that $g^k(\overline{X} - V) \subset U$ and $g^{-k}(\overline{X} - U) \subset V$ whenever $k \geq n$.*

Theorem 2.11 in particular implies $g(+\infty)$ and $g(-\infty)$ are the only fixed points of g in $\overline{X}$ under the conditions of the theorem.

Let G be a group acting by isometries on a Hadamard space X . The action is said to be *proper* if for any compact subset $K \subset X$ the set $\{g \in G : g(K) \cap K \neq \emptyset\}$ is finite. G also acts on $\overline{X}$ as homeomorphisms. A subset $A \subset \overline{X}$ is G -invariant if $g(A) = A$ for all $g \in G$. Theorem 2.11 has the following two corollaries.

Corollary 2.12. *Let X be a Hadamard space that is either locally compact or Gromov hyperbolic, G a group of isometries of X and $g \in G$ a rank one isometry with fixed points $g(+\infty)$, $g(-\infty)$. Then one of the following holds:*

- (i) $g(+\infty)$ or $g(-\infty)$ is fixed by all elements of G ;
- (ii) some axis c of g is G -invariant;
- (iii) G contains a free group of rank two.

Recall a group is *virtually free abelian* if a finite index subgroup is free abelian. A virtually infinite cyclic group is similarly defined.

Corollary 2.13. *Let G act properly and cocompactly by isometries on a $CAT(0)$ space, and H a subgroup of G . If H contains a rank one isometry, then H either is virtually infinite cyclic or contains a free group of rank two.*

For any two isometries g and h of a Hadamard space X , $\langle g, h \rangle$ denotes the group generated by g and h .

Theorem 2.14. ([R]) *Let X be a locally compact Hadamard space and g, h two hyperbolic isometries of X . If $d_T(\xi, \eta) > \pi$ whenever $\xi \in \{g(+\infty), g(-\infty)\}$ and $\eta \in \{h(+\infty), h(-\infty)\}$, then $\langle g, h \rangle$ contains a free group of rank two.*

It follows from Theorem 2.14 that if there are two distinct Tits components C_1 and C_2 such that $\{g(+\infty), g(-\infty)\} \subset C_1$ and $\{h(+\infty), h(-\infty)\} \subset C_2$, then the group $\langle g, h \rangle$ contains a free group of rank two.

2.5. Action on the Limit Set. Let $Isom(X)$ be the group of all isometries of a Hadamard space X , and $\Gamma \subset Isom(X)$ any subgroup. A point $\xi \in \partial_\infty X$ is a *limit point* of Γ if there is a sequence of elements $\{\gamma_i\}_{i=1}^\infty \subset \Gamma$ with $\gamma_i(x) \rightarrow \xi$ for some (hence any) $x \in X$. The *limit set* $\Lambda(\Gamma) \subset \partial_\infty X$ of Γ is the set of limit points of Γ . It is easy to check that $\Lambda(\Gamma)$ is closed (in the cone topology) and Γ -invariant.

Definition 2.15. A nonempty closed and Γ -invariant subset $M \subset \partial_\infty X$ is Γ -*minimal* if it does not contain any proper subset that is closed and Γ -invariant.

When the Hadamard space X is locally compact, the geometric boundary $\partial_\infty X$ and all its closed subsets are compact. It follows from Zorn's lemma that Γ -minimal set always exists when X is locally compact.

We recall two point $\xi, \eta \in \partial_\infty X$ are Γ -*dual* if there is a sequence of elements $\{\gamma_i\}_{i=1}^\infty \subset \Gamma$ such that $\gamma_i(x) \rightarrow \xi$ and $\gamma_i^{-1}(x) \rightarrow \eta$ for some (hence any) $x \in X$ as $i \rightarrow \infty$. Clearly if $\xi, \eta \in \partial_\infty X$ are Γ -dual then $\xi, \eta \in \Lambda(\Gamma)$. For any $\xi \in \Lambda(\Gamma)$, $D_\xi \subset \Lambda(\Gamma)$ denotes the set of points that are Γ -dual to ξ . It is not hard to check that D_ξ is closed and Γ -invariant.

Lemma 2.16. ([B]) *Let X be a locally compact Hadamard space and $\Gamma \subset Isom(X)$ a subgroup. If $\xi, \eta \in \partial_\infty X$ are Γ -dual and $d_T(\xi, \eta) > \pi$, then Γ contains rank one isometries.*

Lemma 2.17. ([CE]) *Let X be a Hadamard space and $\Gamma \subset \text{Isom}(X)$ a subgroup. If $\xi \in \Lambda(\Gamma)$ and $\eta \in \partial_\infty X$ are the endpoints of a geodesic in X , then $D_\xi \subset \overline{\Gamma(\eta)}$.*

The argument in the following proof belongs to Ballmann and Buyalo ([BB]), who assumed $\Lambda(\Gamma) = \partial_\infty X$.

Proposition 2.18. *Suppose Γ is a group of isometries of a locally compact Hadamard space X . If Γ does not contain any rank one isometry, then $d_T(m, \eta) \leq \pi$ for any Γ -minimal set $M \subset \Lambda(\Gamma)$ and any $m \in M$, $\xi \in \Lambda(\Gamma)$.*

Proof. Let $M \subset \Lambda(\Gamma)$ be an arbitrary Γ -minimal set and $m \in M$. Clearly $M = \overline{\Gamma(m)}$. Assume there is some $\eta \in \Lambda(\Gamma)$ with $d_T(m, \eta) > \pi$. By Proposition 2.3(ii), there is a geodesic in X with m and η as endpoints. Lemma 2.17 implies $D_\eta \subset \overline{\Gamma(m)} = M$. Since D_η is nonempty, closed and Γ -invariant and M is Γ -minimal, $D_\eta = M$. In particular, $m \in M = D_\eta$ and m, η are Γ -dual. Now Lemma 2.16 implies there are rank one isometries in Γ since we assumed $d_T(m, \eta) > \pi$. □

Corollary 2.19. *Suppose Γ is a group of isometries of a locally compact Hadamard space X . If Γ does not contain any rank one isometry, then $\Lambda(\Gamma)$ has diameter at most 2π in the Tits metric.*

3. 3-MANIFOLD GROUPS

In this section we study 3-manifold groups and establish the Tits alternative for fundamental groups of closed 3-manifolds with nonpositive sectional curvature (Theorem 3.7). Although Theorem 3.3 should be known to many people, we still include a sketch of the proof. The main reason for doing so is that our proof uses group actions on trees and is a simplified version of the proof in Section 4. The reader is referred to [H] and [K] for definitions and basic facts concerning 3-manifolds.

Definition 3.1. A group G has the *TA-property* if it is virtually free abelian or contains a free group of rank two.

The following lemma is clear.

Lemma 3.2. *A group has the TA-property if some finite index subgroup does.*

Theorem 3.3. *Let M be a Haken 3-manifold whose boundary has zero Euler characteristic, and $H \subset \pi_1(M)$ a subgroup. Then H has the TA-property except in the following two cases:*

- (i) *M is finitely covered by a torus bundle over S^1 and H has finite index in $\pi_1(M)$;*
- (ii) *M is finitely covered by a S^1 -bundle over the torus and H has finite index in $\pi_1(M)$.*

Sketch of proof. By Lemma 3.2 we may assume M is orientable by considering its orientable double cover if necessary. M admits the so-called JSJ decomposition: there is a collection of disjoint embedded tori $\{T_1, \dots, T_k\}$ in M such that the homomorphism $\pi_1(T_i) \rightarrow \pi_1(M)$ induced by inclusion is injective for each i , and each component of $M - \cup_{i=1}^k T_i$ is either a Seifert manifold or an atoroidal manifold. Let $\pi : \tilde{M} \rightarrow M$ be the projection from the universal cover to M . Each component of $\cup \pi^{-1}(T_i)$ is homeomorphic to $\mathbb{R}^2$, and is called a *plane*. Now we construct a graph T from the induced decomposition of $\tilde{M}$. The vertex set of T is in one-to-one correspondence with the set of components of $\tilde{M} - \cup_{i=1}^k \pi^{-1}(T_i)$. Two vertices are joined by an edge if the intersection of the closures of the corresponding components is a plane. The fact that each plane is separating implies the graph T is actually a tree. The action of $\pi_1(M)$ on $\tilde{M}$ preserves the decomposition and induces an action on the tree T . Hence any subgroup $H \subset \pi_1(M)$ also acts on T .

For the action of a group H on a tree T : (see [PV]) if H does not contain a free group of rank two, then one of the following holds: (1) H fixes a point in T , (2) H fixes some $\xi \in \partial_\infty T$, (3) H leaves invariant a geodesic c in T . Now we need to analyze these three cases.

First suppose H fixes a point in T . We may assume H fixes a vertex of T by passing to an index two subgroup if necessary. The component of $\tilde{M} - \cup_{i=1}^k \pi^{-1}(T_i)$ corresponding to this vertex projects down either to a Seifert manifold N or to an atoroidal manifold N , and H is a subgroup of $\pi_1(N)$. If N is an atoroidal manifold, Thurston's theorem says N admits a hyperbolic structure with finite volume. In this case it is clear that H has the TA-property. If N is a Seifert manifold, then H has the TA-property unless H has finite index in $\pi_1(M)$ and M is finitely covered by a circle bundle over the torus.

Suppose H stabilizes a geodesic c in T . After passing to an index two subgroup if necessary, each $h \in H$ translates c . By considering the restricted action of H on c we have an exact sequence: $1 \rightarrow H' \rightarrow H \rightarrow \mathbb{Z} \rightarrow 1$, where $H' \subset H$ is the subgroup consisting of elements of H that fix c pointwise. Since the stabilizer of each edge in $\pi_1(M)$ is $\mathbb{Z}^2$, H' is $\mathbb{Z}^k$ with $k \leq 2$. If $k \leq 1$, then H is clearly virtually $\mathbb{Z}^{k+1}$. If $k = 2$, by consideration of cohomological dimension we see H has finite index in $\pi_1(M)$ and M is finitely covered by a torus bundle over a circle. The case when H fixes some $\xi \in \partial_\infty T$ can be handled similarly. □

Remark 3.4. In the two exceptional cases of Theorem 3.3, both $\pi_1(M)$ and H are virtually solvable.

Our next goal is to establish the Tits alternative for fundamental groups of closed 3-manifolds with nonpositive sectional curvature. The following is a special case of E. Swenson's theorem.

Theorem 3.5. ([Sw]) *Let X be a Hadamard space and G a group acting properly and cocompactly by isometries on X . Suppose $H, K \subset G$ are subgroups and $A, B \subset X$ are closed convex subsets such that $h(A) = A$, $k(B) = B$ for all $h \in H, k \in K$ and $A/H, B/K$ are compact. If $A \cap B \neq \emptyset$, then $H \cap K$ acts cocompactly on $A \cap B$.*

Recall for any group G and any $g \in G$, the centralizer of g in G is $C_g(G) = \{\gamma \in G : \gamma g = g\gamma\}$.

Theorem 3.6. ([R]) *Let X be a Hadamard space, G a group acting properly and cocompactly by isometries on X and $g \in G$ a hyperbolic isometry. Then $\text{Min}(g)$ is invariant under $C_g(G)$ and $C_g(G)$ acts cocompactly on $\text{Min}(g)$.*

Notice an orientable irreducible closed 3-manifold is a Haken manifold if it contains a torus or Klein bottle such that the inclusion induces an injective homomorphism between the fundamental groups.

Theorem 3.7. *Let M be a closed 3-manifold with nonpositive sectional curvature. Then every subgroup $H \subset \pi_1(M)$ has the TA-property.*

Proof. We may assume M is orientable. Notice the theorem follows from Theorem 3.3 if M is a Haken manifold since any solvable subgroup of a group acting properly and cocompactly by isometries on a Hadamard space is virtually free abelian (see [BH]). Let $\pi : X \rightarrow M$ be the universal cover of M . By a theorem of Eberlein (see [E3]) either X contains a 2-flat or $\partial_T X$ is discrete. If $\partial_T X$ is discrete, then each $g \in \pi_1(M)$ is a rank one isometry and the theorem follows easily from Corollary 2.12.

Suppose X contains a 2-flat. Then a theorem of Schroeder ([S2]) says X contains a closed 2-flat F , that is, F is a 2-flat and $\text{Stab}_{\pi_1(M)} F$ acts cocompactly on F . If for any $g \in \pi_1(M)$ either $g(F) = F$ or $g(F) \cap F = \emptyset$ holds, then $\pi(F)$ is an embedded torus or Klein bottle and the inclusion induces an injective homomorphism on the fundamental groups. It follows that M is a Haken manifold.

Now suppose A is a closed 2-flat and there is a $g \in \pi_1(M)$ such that $g(A) \neq A$ and $g(A) \cap A \neq \emptyset$. Since $\dim M = 3$ and A and $g(A)$ are totally geodesic, the intersection $c = A \cap g(A)$ is a complete geodesic. Set $H = \text{Stab}_{\pi_1(M)} A$, $K = gHg^{-1}$ and $B = g(A)$. Then $K = \text{Stab}_{\pi_1(M)} B$ and H, K and A, B satisfy the assumptions in Theorem 3.5. It follows that $H \cap K$ acts cocompactly on $c = A \cap B$ and therefore must be infinite cyclic. Let h be a generator of $H \cap K$ and P_c the parallel set of c . Since $A, B \subset P_c$ and $\dim M = 3$, $P_c = Y \times R$ for a surface Y which is closed and convex in X .

$c = \{y_0\} \times R$ for some $y_0 \in Y$. $h(P_c) = P_c$ and h acts on P_c as $h = (h_0, t)$, where h_0 is an isometry of Y fixing y_0 and t is a translation on R . Since h leaves A invariant, by replacing h with h^2 if necessary we may assume h_0 is trivial. It follows that $P_c = \text{Min}(h)$. By Theorem 3.6 $C_h(\pi_1(M))$ acts cocompactly on $\text{Min}(h) = Y \times R$. If $\partial Y = \phi$ then $X = Y \times R$ and the theorem follows from results on surface groups.

Suppose $\partial Y \neq \phi$. Let $Y_0 \subset Y$ be the convex core of Y , that is, the smallest closed convex subset of Y with $\partial_\infty Y_0 = \partial_\infty Y$ (see [L]). Then $Y \subset N_\epsilon(Y_0)$ for some $\epsilon > 0$ and ∂Y_0 consists of disjoint complete geodesics. Note $C_h(\pi_1(M))$ leaves $Y_0 \times R$ invariant and acts on it cocompactly. Hence there is some $a > 0$ such that $d(c_1, c_2) > a$ for any two distinct geodesics c_1 and c_2 in ∂Y_0 . Fix a geodesic $c_1 \subset \partial Y_0$ and set $F_1 = c_1 \times R$. Now it suffices to show that for any $g \in \pi_1(M)$ either $g(F_1) = F_1$ or $g(F_1) \cap F_1 = \phi$.

Suppose there is some $g \in \pi_1(M)$ with $g(F_1) \neq F_1$ and $g(F_1) \cap F_1 \neq \phi$. Set $F_2 = g(F_1)$. Then $c' = F_1 \cap F_2$ is a complete geodesic. If c' is parallel to c , then $F_2 \subset P_c = \text{Min}(h)$, contradicting to the fact that F_1, F_2 intersect transversally. So c' is not parallel to c . Choose a geodesic $c'' \subset F_2 \cap (Y_0 \times R)$ such that c'' is parallel to c' and $0 < d(c', c'') < a$. Then $c'' \subset \gamma \times R$ for a complete geodesic $\gamma \subset Y_0$. Since c'' and c' are parallel, γ and c_1 are parallel and bound a flat strip in Y_0 . The fact that Y_0 is the convex core of Y implies $\gamma = c_1$ and so $c'' \subset F_1$, contradicting to the fact that F_1, F_2 intersect transversally. □

4. DECOMPOSITION OF REAL ANALYTIC 4-MANIFOLDS

In this section we shall prove one of the main results (Theorem 4.1) of the paper. The definition of a cycle is given in Definition 2.8.

Theorem 4.1. *Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature. Suppose there are no cycles in X . If a subgroup of Γ is not virtually free abelian, then it contains a free group of rank two.*

Remark 4.2. *There exist real analytic closed 4-manifolds (see [AS]) satisfying the assumptions of the theorem.*

Let $M = X/\Gamma$ be as in Theorem 4.1. We shall decompose X into convex domains and construct a 2-complex Y associated to the decomposition. The 2-complex Y is a $CAT(-1)$ space and the group Γ acts on Y as a group of isometries. Thus any subgroup H of Γ also acts on Y as isometries. A group acting on a $CAT(-1)$ space contains a free group of rank two unless it fixes a point in $\bar{Y} = Y \cup \partial_\infty Y$ or stabilizes a geodesic. Therefore it suffices to consider these special cases.

The decomposition and the associated 2-complex should be considered as analogues of JSJ decomposition and the associated Bass-Serre tree in 3-manifold theory.

4.1. Decomposition and the Associated 2-complex. Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature. In this section we shall decompose X , construct the associated 2-complex Y and prove Y is a $CAT(-1)$ space.

We use the results and notation of Section 2.3. Recall $\mathcal{W}$ is the set of maximal higher rank submanifolds of X that are of the form $Q \times R$ for nonflat 2-dimensional Hadamard manifolds Q . Each $W \in \mathcal{W}$ is called a *wall*. We notice each wall W is totally geodesic and $X - W$ has two components. Set $X_0 = X - \cup_{W \in \mathcal{W}} W$. We say two components C_1 and C_2 of X_0 are separated by a wall W if they lie in different components of $X - W$, and say C_1 and C_2 lie on the same side of W if they lie in the same component of $X - W$. Similarly we define two points $x, y \in X - W$ either to be separated by W or to lie on the same side of W .

Suppose W_1, W_2 are two walls with $W_1 \cap W_2 \neq \phi$. Then $W_1 \cap W_2 \cap W = \phi$ for any wall $W \neq W_1, W_2$. Since each wall is closed, and there are only a finite number of walls modulo Γ , there is some $\epsilon > 0$ depending only on M with $d(W_1 \cap W_2, W) > \epsilon$ for any wall $W \neq W_1, W_2$. It follows that

there are 4 components C_1, C_2, C_3, C_4 of X_0 with $W_1 \cap W_2 \subset \overline{C}_i$ ($i = 1, 2, 3, 4$) and $W_1 \cap W_2 \cap \overline{C} = \emptyset$ for any component C of X_0 , $C \neq C_1, C_2, C_3, C_4$. By suitably labeling the 4 components C_1, C_2, C_3, C_4 , we may assume the following intersections are 3-dimensional: $\overline{C}_1 \cap \overline{C}_2 \subset W_1$, $\overline{C}_2 \cap \overline{C}_3 \subset W_2$, $\overline{C}_3 \cap \overline{C}_4 \subset W_1$, $\overline{C}_4 \cap \overline{C}_1 \subset W_2$.

Now we construct the 2-complex Y . The set V of vertices of Y is in one-to-one correspondence with the set of components of X_0 . We denote by C_v the component of X_0 that corresponds to the vertex $v \in V$. Two vertices v_1 and v_2 are joined by an edge (denoted by $v_1 v_2$) if and only if $\overline{C}_{v_1} \cap \overline{C}_{v_2}$ is three dimensional. When W_1, W_2 are two walls with $W_1 \cap W_2 \neq \emptyset$, there are 4 components $C_{v_1}, C_{v_2}, C_{v_3}, C_{v_4}$ of X_0 such that the following intersections are 3-dimensional: $\overline{C}_{v_1} \cap \overline{C}_{v_2} \subset W_1$, $\overline{C}_{v_2} \cap \overline{C}_{v_3} \subset W_2$, $\overline{C}_{v_3} \cap \overline{C}_{v_4} \subset W_1$, $\overline{C}_{v_4} \cap \overline{C}_{v_1} \subset W_2$. Thus there are the following edges in Y : $v_1 v_2, v_2 v_3, v_3 v_4, v_4 v_1$. Now whenever there are two walls W_1, W_2 with $W_1 \cap W_2 \neq \emptyset$, and $v_1 v_2, v_2 v_3, v_3 v_4, v_4 v_1$ the corresponding edges in Y as described above, we attach a square (denoted by $S(W_1, W_2)$) along these 4 edges. The construction of Y is complete.

Next we will put a metric on Y . A *hyperbolic square* is a closed convex region in the real hyperbolic plane whose boundary is the union of 4 geodesic segments such that the 4 geodesic segments are of the same length and the interior angles at the endpoints of these geodesic segments are all equal; each of the 4 geodesic segments is called an edge and the endpoints of the 4 geodesic segments are called vertices. Let S_0 be a hyperbolic square so that the interior angle α at the vertices satisfies: $\alpha > \frac{2}{5}\pi$. Let l be the length of an edge of S_0 . Now we declare that all the edges in Y have length l , and all the squares in Y are isometric to S_0 . Thus Y is a piecewise hyperbolic 2-complex. Since there are only a finite number (actually 3 types) of isometry types of cells in Y , Y with the path metric is a complete geodesic metric space. We shall show that Y is a $CAT(-1)$ space.

We first look at the intersection $\overline{C}_{v_1} \cap \overline{C}_{v_2}$ for each edge $v_1 v_2$ of Y .

Lemma 4.3. *Let $W = Q \times R$ be a wall so that $W \cap W_1 \neq \emptyset$ for some wall $W_1 \neq W$. Suppose Z is a component of $W - \cup_{W' \neq W} W'$, where W' varies over all walls that are distinct from W . Then the closure $\overline{Z}$ of Z has the form: $\overline{Z} = Q' \times R \subset Q \times R = W$, where $Q' \subset Q$ is a closed convex subset of Q and is the universal cover of a nonpositively curved compact surface with closed geodesics on the boundary.*

Proof. Recall (see Theorem 2.6) if $W \cap W' \neq \emptyset$, then $W \cap W' = c \times R \subset Q \times R$ for a complete geodesic c in Q . And $W \cap W' \cap W'' = \emptyset$ if W, W', W'' are three distinct walls. Thus $W - \cup_{W' \neq W} W' = Q \times R - \cup_i (c_i \times R)$, where $\{c_i\}$ is a disjoint collection of complete geodesics in Q . Since W is closed and there are only a finite number of maximal higher rank submanifolds modulo Γ , the lemma follows. $\square$

For any wall W , let $X_W = X - \cup_{W' \neq W} W'$ where W' varies over all walls different from W .

Lemma 4.4. *Let $v_1 v_2$ be an edge of Y . Then:*

- (i) C_{v_1}, C_{v_2} are separated by a unique wall W ;
- (ii) for any $x_1 \in C_{v_1}$ and $x_2 \in C_{v_2}$, W is the only wall that intersects the geodesic segment $x_1 x_2$, and the intersection is transversal;
- (iii) if E is the component of X_W that contains C_{v_1} , then C_{v_1} and C_{v_2} are the only two components of X_0 contained in E .

Proof. (i) Since v_1, v_2 are distinct, C_{v_1}, C_{v_2} are distinct and are thus separated by at least one wall. By the construction of Y , $\overline{C}_{v_1} \cap \overline{C}_{v_2}$ is three dimensional. Let W be any wall that separates C_{v_1} and C_{v_2} . Then $\overline{C}_{v_1} \cap \overline{C}_{v_2} \subset W^+ \cap W^- = W$, where W^+ and W^- are the closures of the two components of $X - W$. Since for any two distinct walls W_1, W_2 the intersection $W_1 \cap W_2$ is at most 2-dimensional, there is exactly one wall W that separates C_{v_1} and C_{v_2} .

(ii) Let W' be a wall with $W' \cap x_1 x_2 \neq \emptyset$. The intersection must be transversal since W' is totally geodesic and $x_1 \notin W'$. Thus x_1 and x_2 are in different components of $X - W'$. It follows

that C_{v_1} and C_{v_2} are contained in different components of $X - W'$ and W' separates C_{v_1} and C_{v_2} . Now (i) implies $W' = W$.

(iii) From (ii) we see for any $x_1 \in C_{v_1}$ and $x_2 \in C_{v_2}$, the segment x_1x_2 is contained in E . Thus E contains both C_{v_1} and C_{v_2} . Let $C \neq C_{v_1}$ be a component of X_0 contained in E . $C \neq C_{v_1}$ implies C and C_{v_1} are separated by a wall, which must be W by the definition of E . Now there is no wall separating C_{v_2} and C and we have $C = C_{v_2}$.

□

By the Cartan-Hadamard theorem, to prove Y is $CAT(-1)$ it suffices to show Y is simply connected and has curvature ≤ -1 . Since Y is piecewise hyperbolic, Y has curvature ≤ -1 if all the vertex links are $CAT(1)$. The vertex links are metric graphs and a metric graph is $CAT(1)$ if and only if each injective edge loop has length at least 2π . By the construction of Y , the edges in the vertex links all have the same length $\alpha > \frac{2}{5}\pi$. Therefore the vertex links are $CAT(1)$ if each injective edge loop (in the links) has at least 5 edges.

Lemma 4.5. *Let $v \in Y$ be a vertex of Y and $Link(v, Y)$ the link of v in Y . Then each injective edge loop in $Link(v, Y)$ has at least 5 edges. In particular, Y has curvature ≤ -1 .*

Proof. Let e be an edge in $Link(v, Y)$. Then e corresponds to a square S of Y that has v as one of its 4 vertices. Let v, v_1, v', v_2 be the 4 vertices of S in cyclic order. By Lemma 4.4, there is a unique wall W_1 that separates C_v and C_{v_1} and a unique wall W_2 that separates C_v and C_{v_2} . By the construction of Y , $W_1 \cap W_2 \neq \emptyset$ and the square S is determined by W_1 and W_2 . The two endpoints of e uniquely determine v_1 and v_2 , v_1 and v_2 then uniquely determine W_1 and W_2 , and W_1 and W_2 in turn uniquely determine the square S . It follows that the edge e is uniquely determined by its two endpoints. Therefore there is no injective edge loop with length 2 in $Link(v, Y)$.

Now let L be an injective edge loop in $Link(v, Y)$ consisting of n edges. Then there are n edges vv_i ($i = 1, 2, \dots, n$) in Y so that vv_i, vv_{i+1} (indices are taken modulo n) are two edges of a square. It follows from the preceding paragraph that there are walls $W_1, \dots, W_n$ such that W_i is the only wall separating C_v and C_{v_i} and $W_i \cap W_{i+1} \neq \emptyset$ for all i where indices are taken modulo n . Suppose there are indices i and j with $i \neq j$ and $W_i = W_j$. Let E be the component of X_{W_i} containing C_v . Lemma 4.4 applied to the edge vv_i implies $C_v, C_{v_i} \subset E$. Similarly $C_v, C_{v_j} \subset E$ as $W_i = W_j$. Since L is an injective loop in $Link(v, Y)$, $i \neq j$ implies $v_i \neq v_j$. Therefore $C_{v_i} \neq C_{v_j}$ and there are three distinct components C_v, C_{v_i}, C_{v_j} of X_0 contained in E , contradicting to Lemma 4.4. It follows that $W_1, \dots, W_n$ are all distinct and form an n -cycle in X . Now Proposition 2.9 completes the proof.

□

Lemma 4.6. *Any edge loop in Y is homotopically trivial. In particular, Y is simply connected.*

Proof. We induct on the length of an edge loop. By the construction of Y , two vertices are connected by at most one edge and there is no injective edge loop with length 2.

Let l be an injective edge loop in Y and $v_1, \dots, v_n$ the vertices on l in cyclic order. Let $x_i \in C_{v_i}$ be an arbitrary point. Notice the components C_{v_i} ($i = 1, \dots, n$) are all distinct since l is an injective edge loop. Denote by W_i the unique wall that separates C_{v_i} and $C_{v_{i+1}}$ (here indices are taken modulo n). By Lemma 4.4 W_i is the only wall intersecting $x_i x_{i+1}$. Let $L = \cup_{i=1}^n x_i x_{i+1}$. Clearly L is a loop in X .

Suppose $n = 3$. Since W_1 separates x_1 and x_2 , the path $x_2 x_3 * x_3 x_1$ must intersect W_1 . From the preceding paragraph the path $x_2 x_3 * x_3 x_1$ only intersects W_2 and W_3 , we have $W_2 = W_1$ or $W_3 = W_1$. We may assume $W_2 = W_1$, the other case being handled similarly. Let E_2 be the component of X_{W_2} that contains C_{v_2} . Then Lemma 4.4 implies the three distinct components $C_{v_1}, C_{v_2}, C_{v_3}$ are all contained in E_2 since $W_2 = W_1$, which contradicts to the same lemma. It follows that there is no injective edge loop with length 3.

Now suppose $n \geq 4$. Since W_1 separates x_1 and x_2 , the path $L - x_1x_2$ must intersect W_1 . By construction the path $L - x_1x_2$ only intersects $W_2, \dots, W_n$. So $W_i = W_1$ for at least one i , $2 \leq i \leq n$. Let m be the largest such i . Then x_1 and x_{m+1} lie on one side of W_1 , and x_2 and x_m lie on the other side of W_1 . The argument in the preceding paragraph shows $x_1 \neq x_{m+1}$ and $x_2 \neq x_m$ since l is an injective loop. Let $z = x_1x_2 \cap W_1$ and $z' = x_mx_{m+1} \cap W_1$. Since $x_2 \neq x_m$, we have $C_{v_m} \neq C_{v_2}$, and by Lemma 4.4 (iii) there is at least one wall W' , $W' \neq W_1$ with $W' \cap zz' \neq \emptyset$. Let $\{W'_1, \dots, W'_k\}$ be the set of walls different from W_1 that intersect zz' . Then $W_1 \cap W'_i \neq \emptyset$ ($i = 1, \dots, k$) and $S(W_1, W'_i)$ is a square in Y . Set $z_i = zz' \cap W'_i$. We label the walls $W'_1, \dots, W'_k$ so that $d(z, z_i) < d(z, z_j)$ whenever $i < j$. Then the interior of zz_1 does not intersect any wall different from W_1 . It follows that v_1, v_2 are vertices of $S(W_1, W'_1)$. Let v'_2, v''_2 be the other two vertices of $S(W_1, W'_1)$ so that $C_{v'_2}, C_{v''_2}$ lie on the same side of W_1 . Similarly we define v'_i, v''_i for all $2 \leq i \leq k+1$ so that $v'_{k+1} = v_m, v''_{k+1} = v_{m+1}$ and all $C_{v'_i}$ ($2 \leq i \leq k+1$) lie on the same side of W_1 .

Let $l_1 = v_2v'_2 \cdots v'_kv_m, l_2 = v_1v''_2 \cdots v''_kv_{m+1}, l_3 = v_2v_3 \cdots v_{m-1}v_m, l_4 = v_{m+1}v_{m+2} \cdots v_nv_1$ be oriented paths in Y and l_i^{-1} ($i = 1, 2, 3, 4$) the same paths with the reverse orientation. Set $l' = v_1v_2 * l_3 * l_1^{-1} * v_2v_1, l'' = v_1v_2 * l_1 * v_mv_{m+1} * l_2^{-1}, l''' = l_2 * l_4$. Then l', l'' and l''' are oriented loops so that $l = v_1v_2 * l_3 * v_mv_{m+1} * l_4$ is homotopic to $l' * l'' * l'''$. Notice $\cup_{i=1}^k S(W_1, W'_i)$ is homeomorphic to a square and l'' is its boundary. Therefore l'' is homotopically trivial. Now we notice $\text{length}(l_3) \geq k$ since the path $x_2x_3 * \cdots * x_{m-1}x_m$ must cross all the walls $W'_1, \dots, W'_k$. Similarly $\text{length}(l_4) \geq k$. Since $\text{length}(l_1) = \text{length}(l_2) = k$, the lengths of the loops $l_3 * l_1^{-1}$ and $l_2 * l_4$ are strictly less than the length of l . The induction hypothesis implies $l_3 * l_1^{-1}$ and $l_2 * l_4$ are homotopically trivial. Therefore l' and l''' are homotopically trivial. It follows l is also homotopically trivial. $\square$

Lemmas 4.5 and 4.6 together imply the following proposition.

Proposition 4.7. The 2-complex Y is a $CAT(-1)$ space.

Recall a group is said to have the TA-property if it is virtually free abelian or contains a free group of rank two. We note Theorem 4.1 holds if and only if any subgroup H of Γ has the TA-property. Let $H \subset \Gamma$ be a subgroup of Γ . By Lemma 3.2 we may assume each $h \in H$ preserves the orientation of X , after passing to an index two subgroup if necessary. Corollary 2.13 implies we may assume H does not contain any rank one isometry. By Theorem 2.14, we may further assume there is a Tits component $\mathcal{C} \subset \partial_T X$ so that $h(+\infty), h(-\infty) \in \mathcal{C}$ for all $h \in H$. If $\mathcal{C} = \partial_T F$ for a maximal higher rank submanifold F which is a 2-flat or 3-flat, then each element $h \in H$ leaves F invariant. It follows that H acts on the Euclidean space F properly and isometrically, and thus must be virtually free abelian by Bieberbach's theorem.

From now on we assume $H \subset \Gamma$ satisfies the following properties:

- (a) each $h \in H$ preserves the orientation of X ;
- (b) H does not contain any rank one isometry;
- (c) there is a unique Tits component $\mathcal{C} \subset \partial_T X$ such that $h(+\infty), h(-\infty) \in \mathcal{C}$ for all $h \in H$;
- (d) the Tits component $\mathcal{C}$ has the following form: $\mathcal{C} = \bigcup_{W \in \mathcal{W}^*} \partial_\infty W$, where $\mathcal{W}^* \subset \mathcal{W}$ is a subset such that $\bigcup_{W \in \mathcal{W}^*} W$ is a connected component of $\bigcup_{W \in \mathcal{W}} W \subset X$.

We shall also frequently pass to finite index subgroups of H .

4.2. Action on the 2-complex. Since $\mathcal{W}$ is invariant under the action of the group Γ , it is clear from the construction of Y that Γ acts on Y as a group of cellular isometries. Any cellular isometry of Y either is hyperbolic or have a fixed point in Y as Y only has a finite number of isometry types of cells (see [Br]). The fact that Y is a $CAT(-1)$ space implies any hyperbolic isometry of Y is of rank one. Thus any $\gamma \in \Gamma$ either acts on Y as a rank one isometry or has a fixed point in Y . Notice

an element $\gamma \in \Gamma$ may act as a rank one isometry on Y even if it is not rank one (with respect to its action on X) in Γ .

For any $\gamma \in \Gamma$, let $\gamma_Y : Y \rightarrow Y$ denote the induced isometry of γ on Y , and $\text{Fix}(\gamma_Y) \subset Y$ denote the fixed point set of γ_Y . Let $H \subset \Gamma$ be a subgroup satisfying the properties stated at the end of Section 4.1. As a subgroup of Γ , H also acts on Y as isometries. We shall analyze the action of H and its individual elements on Y .

Lemma 4.8. *Let $S = S(W_1, W_2)$ be a square in Y and $\gamma \in \Gamma$. If $\gamma_Y(S) = S$, then $\gamma(W_1 \cap W_2) = W_1 \cap W_2$.*

Proof. Since W_1, W_2 are uniquely determined by S , $\gamma_Y(S) = S$ implies $\{\gamma(W_1), \gamma(W_2)\} = \{W_1, W_2\}$. The lemma follows. $\square$

Let $e = v_1 v_2$ be an edge in Y , and C_1, C_2 the two components of X_0 corresponding to v_1, v_2 respectively. Lemma 4.4 implies there is a unique wall separating C_1 and C_2 . The following lemma is clear.

Lemma 4.9. *Let $e = v_1 v_2$ be an edge in Y , C_1, C_2 the two components of X_0 corresponding to v_1, v_2 respectively, and W the unique wall separating C_1 and C_2 . If $\gamma_Y(e) = e$ for some $\gamma \in \Gamma$, then $\gamma(W) = W$.*

Suppose $\gamma \in \Gamma$ is not a rank one isometry. Let $P(\gamma)$ denote the parallel set of an axis of γ . Since X is real analytic, $P(\gamma)$ is a higher rank submanifold in X , and therefore is contained in a maximal higher rank submanifold. We further suppose $P(\gamma) \subset W$ for a wall $W = Q \times R$, where Q is a 2-dimensional nonflat Hadamard manifold. Notice γ belongs to exactly one of the following 4 classes:

Type A: $P(\gamma) = c \times R \subset Q \times R = W$ and $P(\gamma) \cap W' = \emptyset$ for any wall $W' \neq W$, where c is a complete geodesic in Q ;

Type B: $P(\gamma) = W \cap W'$ for some wall $W' \neq W$;

Type C: $P(\gamma) = W$. In this case, $\{q_0\} \times R$ is an axis of γ for some $q_0 \in Q$;

Type D: $P(\gamma) = c \times R \subset Q \times R = W$ and the axes of γ intersect transversally with some wall W' , where c is a complete geodesic in Q .

Before we study the action of individual isometries on Y , we introduce a subcomplex of Y associated to each wall that intersects other walls. Recall for each edge $v_1 v_2$ in Y there is a unique wall $\tilde{W}$ separating C_{v_1} and C_{v_2} . Given any square $S(W_1, W_2)$ in Y , two opposite edges of $S(W_1, W_2)$ determine W_1 , while the other two opposite edges determine W_2 . Let $\omega(W_1, W_2) \subset S(W_1, W_2)$ be the geodesic segment connecting the midpoints of the two opposite edges of $S(W_1, W_2)$ that determine W_1 . Let W be a wall intersecting other walls. Set $C_W = \bigcup S(W, W')$ and $T_W = \bigcup \omega(W, W') \subset C_W$ where W' varies over all walls $W' \neq W$ with $W' \cap W \neq \emptyset$. C_W is a subcomplex of Y .

Lemma 4.10. *Let $\gamma \in \Gamma$ be a Type C isometry. If the wall $W := P(\gamma)$ intersects other walls, then there is an integer n such that $\text{Min}(\gamma^n) = W$ and $\text{Fix}(\gamma_Y^n) = C_W$. In particular C_W is a closed convex subset of Y , and $\phi \neq \text{Fix}(\gamma_Y) \subset C_W$.*

Proof. Let $W = Q \times R$. By the definition of a Type C isometry, there is some $q_0 \in Q$ such that $\{q_0\} \times R$ is an axis of γ . It follows that $\gamma|_W$ has the following form $\gamma|_W = (e, t) : Q \times R \rightarrow Q \times R$ where e is an elliptic isometry of Q fixing the point q_0 and t is a translation of R . Since the wall W is closed and Q is a nonflat 2-dimensional Hadamard manifold, e has finite order (see [E1]). Therefore there is an even integer n such that $e^n = \text{id}$. Notice $\text{Min}(\gamma^n) = W$ and γ^n preserves the orientation of X .

Now assume $\gamma \in \Gamma$ is a Type C isometry preserving the orientation of X with $\text{Min}(\gamma) = W$. We shall show $\text{Fix}(\gamma_Y) = C_W$. We first argue $C_W \subset \text{Fix}(\gamma_Y)$. Let $W' \neq W$ with $W' \cap W \neq \emptyset$. Since $\text{Min}(\gamma) = W = Q \times R$, each geodesic $\{q\} \times R$ ($q \in Q$) is an axis of γ . It follows that the

2-flat $W \cap W' = c \times R \subset Q \times R = W$ is invariant under γ . Since W' is perpendicular to W , we have $\gamma(W') = W'$. Therefore $\gamma_Y(S(W, W')) = S(W, W')$. Since γ preserves the orientation of X and translates all the geodesics $\{q\} \times R$ ($q \in Q$), each of the 4 components of X_0 determined by W and W' is invariant under γ . So the square $S(W, W')$ is pointwise fixed by γ_Y .

Next we show $\text{Fix}(\gamma_Y) \subset C_W$. Let $y \in \text{Fix}(\gamma_Y)$. If y lies in the interior of a square $S(W_1, W_2)$, then $\gamma_Y(S(W_1, W_2)) = S(W_1, W_2)$. Lemma 4.8 implies $\gamma(W_1 \cap W_2) = W_1 \cap W_2$ and thus an axis of γ lies in $W_1 \cap W_2$. It follows that $W \cap W_1 \cap W_2 \neq \emptyset$. By Theorem 2.5 (iv) $W = W_1$ or $W = W_2$. In either case $S(W_1, W_2) \subset C_W$.

Suppose y lies in the interior of an edge $v_1 v_2$. Then $\gamma_Y(v_1 v_2) = v_1 v_2$. Let W' be the unique wall separating C_{v_1} and C_{v_2} . Lemma 4.9 implies $\gamma(W') = W'$ and thus W' contains an axis of γ . It follows $W \cap W' \neq \emptyset$. If $W' = W$, then clearly $v_1 v_2 \subset C_W$. Suppose $W' \neq W$. Pick $x_1 \in C_{v_1}$, $x_2 \in C_{v_2}$ and let $z' = x_1 x_2 \cap W'$. Also pick $z \in W' \cap W$. Then $zz' \subset W'$. Let $W = W_1, \dots, W_k$ be the sequence of walls in consecutive order that intersect zz' transversally. These walls determine a sequence of squares $S(W', W_1), \dots, S(W', W_k)$ in Y , where two consecutive squares share exactly an edge. Notice $v_1 v_2$ is an edge of $S(W', W_k)$. For each of these squares, two opposite edges correspond to the wall W' . Let l_i ($i = 1, 2, \dots, k$) be the geodesic segment connecting the midpoints of these two opposite edges, and $l = \cup_{i=1}^k l_i$. l is a geodesic by the geometry of Y . One endpoint of l lies in $S(W', W) \subset C_W$ and the other one is the midpoint of the edge $v_1 v_2$, so they are both fixed by γ_Y . It follows that l is pointwise fixed by γ_Y , and by the preceding paragraph $S(W', W_i) \subset C_W$ for each i . In particular $v_1 v_2 \subset C_W$.

Now suppose y is a vertex. Then for any $y' \in C_W$, the geodesic segment yy' is pointwise fixed by γ_Y . The initial segment of yy' is contained in some square or edge. It follows from the above that this square or edge lies in C_W and so does y .

□

It follows from Lemma 4.10 that C_W is a $CAT(-1)$ space. C_W clearly admits a “reflection” about T_W with T_W as the fixed point set. Thus T_W is also a closed and convex subset of Y . T_W is the “core” of C_W , and C_W is homeomorphic to $T_W \times [0, 1]$. For each wall W that intersects other walls, let $T_W(\infty) \subset \partial_\infty Y$ be the geometric boundary of T_W naturally identified with a subset of $\partial_\infty Y$.

Lemma 4.11. *Let $W_1 \neq W_2$ be two walls that intersect other walls. Then $T_{W_1}(\infty) \cap T_{W_2}(\infty) = \emptyset$.*

Proof. Suppose $T_{W_1}(\infty) \cap T_{W_2}(\infty) \neq \emptyset$ and pick $\xi \in T_{W_1}(\infty) \cap T_{W_2}(\infty)$. Let $\alpha_1 : [0, \infty) \rightarrow T_{W_1}$ and $\alpha_2 : [0, \infty) \rightarrow T_{W_2}$ be two rays ending at ξ . Since Y is a $CAT(-1)$ space, we have $d(\alpha_1(t), \alpha_2(t)) \rightarrow 0$ as $t \rightarrow \infty$. Since α_1 and α_2 cross centers of squares in Y , there are $a > 0$, $b \in \mathbb{R}$ so that $\alpha_1(t) = \alpha_2(t + b)$ for all $t \geq a$. α_1 passes through midpoints of edges in Y and such a midpoint uniquely determines the wall W_1 . The same is true for α_2 and W_2 . Hence $W_1 = W_2$, a contradiction.

□

Let γ be a Type A isometry and W the wall with $P(\gamma) \subset W$. Then $P(\gamma) \subset X_W$. Let E be the component of X_W containing $P(\gamma)$, and C_1 and C_2 the two components of X_0 contained in E . Denote by v_1 and v_2 the two vertices of Y corresponding to C_1 and C_2 respectively. Then $v_1 v_2$ is an edge in Y .

Lemma 4.12. *Let γ be a Type A isometry, and $W, v_1 v_2$ be as above. Then $\phi \neq \text{Fix}(\gamma_Y) \subset v_1 v_2$.*

Proof. We use the notation from the paragraph preceding the lemma. Since E is the component of X_W containing $P(\gamma)$ and $P(\gamma)$ is invariant under γ , $\gamma(E) = E$. It follows that $\{\gamma(C_1), \gamma(C_2)\} = \{C_1, C_2\}$ and γ_Y sends $v_1 v_2$ to itself. In particular, the midpoint of $v_1 v_2$ is fixed by γ_Y . Lemma 4.8 implies no square in Y is invariant under γ_Y . Then Lemma 4.9 and an argument similar to the proof of Lemma 4.10 shows $\text{Fix}(\gamma_Y) \subset v_1 v_2$.

□

Lemma 4.13. *Let γ be a Type B isometry and W_1, W_2 the two walls such that $P(\gamma) = W_1 \cap W_2$. Then $\phi \neq \text{Fix}(\gamma_Y) \subset S(W_1, W_2)$.*

Proof. Since $P(\gamma) = W_1 \cap W_2$, we have $\{\gamma(W_1), \gamma(W_2)\} = \{W_1, W_2\}$ so the square $S(W_1, W_2)$ is invariant under γ_Y . In particular, the center of the square $S(W_1, W_2)$ is fixed by γ_Y . Now an argument similar to the proof of Lemma 4.10 shows $\text{Fix}(\gamma_Y)$ is contained in $S(W_1, W_2)$. $\square$

Lemma 4.14. *Let γ be a Type D isometry with $P(\gamma)$ contained in a wall W . Then γ_Y is a rank one isometry of Y , and the axis of γ_Y is contained in T_W .*

Proof. We exhibit a geodesic in T_W that is translated by γ_Y . Let $c \subset P(\gamma) \subset W$ be an axis of γ . Then c intersects some wall W' transversally since γ is a Type D isometry. Pick a point $p \in c$ such that W is the only wall containing p . Let $W_1, W_2, \dots, W_k$ be the sequence of walls in consecutive order intersected transversally by the geodesic segment $p\gamma(p)$. Then the biinfinite sequence of walls

$$\dots, \gamma^{-1}(W_1), \dots, \gamma^{-1}(W_k), W_1, \dots, W_k, \gamma(W_1), \dots, \gamma(W_k), \dots$$

is the sequence of walls in consecutive order intersected transversally by the axis c . This sequence of walls together with W determine a sequence of squares in Y . The union of this sequence of squares is a “strip” R contained in C_W , and $\gamma_Y(R) = R$. It follows that the geodesic $R \cap T_W$, the center line of R , is translated by γ_Y . $\square$

Let $H \subset \Gamma$ be a subgroup satisfying the properties stated at the end of Section 4.1. As a subgroup of Γ , H also acts on Y as isometries. Y is a Gromov hyperbolic space since it is $CAT(-1)$. By a theorem of Gromov (see [G]), one of the following occurs:

- (1) H has a bounded orbit in Y ;
- (2) H has a fixed point in $\partial_\infty Y$;
- (3) h_Y is a rank one isometry for some $h \in H$.

When (1) occurs, Cartan’s fixed point theorem implies H has a fixed point in Y . After passing to a finite index subgroup, we may assume that H fixes a vertex of Y . In Section 4.3 we show H has the TA-property if it fixes a vertex of Y and there is no cycle in X .

Lemma 4.15. *Suppose H has a fixed point in $\partial_\infty Y$ and h_Y is not a rank one isometry for any $h \in H$. Then H has the TA property.*

Proof. By assumption and the analysis of the 4 types of isometries, each $h \in H$ is of Type C. By Lemma 4.10 if h is of Type C and W is the wall with $P(h) = W$, then the fixed point set of h_Y in $\partial_\infty Y$ is contained in $T_W(\infty)$. Now Lemma 4.11 and the fact that H has a fixed point in $\partial_\infty Y$ imply that there is a wall W with $P(h) = W$ for all $h \in H$. Therefore $h(W) = W$ for all $h \in H$ and $H \subset \text{Stab}_\Gamma W$. Since $\text{Stab}_\Gamma W$ is the fundamental group of the closed 3-manifold $W/\text{Stab}_\Gamma W$ with nonpositive sectional curvature, the lemma follows from Theorem 3.7. $\square$

Now we remain to consider the case when g_Y is rank one for some $g \in H$. Let $c \subset Y$ be the axis of g_Y , and $c(+\infty)$, $c(-\infty)$ the fixed points of g_Y in $\partial_\infty Y$. By Corollary 2.12 one of the following holds:

- (1) $c(+\infty)$ or $c(-\infty)$ is fixed by all elements of H ;
- (2) $h_Y(c) = c$ for all $h \in H$;
- (3) H contains a free group of rank two.

We only need to consider the first two cases. Case (2) can be reduced to case (1) as in case (2) an index two subgroup of H fixes both $c(+\infty)$ and $c(-\infty)$.

Lemma 4.16. *Suppose g_Y is rank one for some $g \in H$, and $c(+\infty)$, $c(-\infty)$ are the fixed points of g_Y in $\partial_\infty Y$. If $c(+\infty)$ or $c(-\infty)$ is fixed by all elements of H , then H has the TA-property.*

Proof. The assumption implies g is of Type D. Let W be the wall with $\text{Min}(g) \subset W$. Then the axis c of g_Y is contained in T_W . Since each $h \in H$ fixes a point in $\partial_\infty Y$, by the analysis of the 4 types of isometries we see each $h \in H$ is either of Type D or of Type C. By Lemmas 4.10, 4.11, 4.14 and the fact that $c(+\infty)$ or $c(-\infty)$ is fixed by all elements of H , we conclude $h(W) = W$ for all $h \in H$. Now the argument in the proof of Lemma 4.15 shows H has the TA-property. $\square$

4.3. Subgroups With a Fixed Point in the 2-complex. We remain to consider the case when H fixes a vertex of Y . To be more precise, throughout this section, $H \subset \Gamma$ is a subgroup satisfying the following properties:

- (a) there is a Tits component $\mathcal{C} \subset \partial_T X$ so that $h(+\infty), h(-\infty) \in \mathcal{C}$ for all $h \in H$;
- (b) $\mathcal{C} = \bigcup_{W \in \mathcal{W}^*} \partial_\infty W$, where $\mathcal{W}^* \subset \mathcal{W}$ is a subset such that $\bigcup_{W \in \mathcal{W}^*} W$ is a connected component of $\bigcup_{W \in \mathcal{W}} W \subset X$;
- (c) there is a vertex v in Y so that $h_Y(v) = v$ for all $h \in H$.

Notice condition (c) implies H contains no Type D isometries since by Lemma 4.14 γ_Y has no fixed point in Y if γ is of Type D.

Set $W_{\mathcal{C}} = \bigcup_{W \in \mathcal{W}^*} W$. Then both C_v and $W_{\mathcal{C}}$ are invariant under H . It follows that $\overline{C}_v \cap W_{\mathcal{C}}$ is invariant under H . Set $\Lambda = \overline{C}_v \cap W_{\mathcal{C}}$. Then $H \subset \text{Stab}_\Gamma \Lambda$. We shall prove Λ is connected and $\Lambda / \text{Stab}_\Gamma \Lambda$ is a compact 3-manifold. Under the assumption that there is no cycle in X , Λ is simply connected and consequently H is a subgroup of the fundamental group of a compact 3-manifold. Theorem 3.3 then implies H has the TA-property.

For any wall W , the closures of the two components of $X - W$ are called closed half spaces. $\overline{C}_v$ is clearly the intersection of a family of closed half spaces. Now it is not hard to see that if W is a wall with $W \cap \overline{C}_v \neq \emptyset$, then $W \cap \overline{C}_v = \overline{Z}$ for a component Z of $W - \bigcup_{W' \neq W} W'$, where W' varies over all walls distinct from W . By Lemma 4.3 such a $\overline{Z}$ has the form $\overline{Z} = Q' \times R \subset Q \times R = W$, where $Q' \subset Q$ is a closed convex subset of Q and is the universal cover of a nonpositively curved compact surface with closed geodesics on the boundary. Similarly we see if $W_1 \neq W_2$ and $W_1 \cap W_2 \cap \overline{C}_v \neq \emptyset$, then $W_1 \cap W_2 \cap \overline{C}_v$ is a 2-flat.

Let $W, W' \in \mathcal{W}^*$. A *chain* from W to W' is a sequence $W = W_0, W_1, \dots, W_n = W'$ of walls in $\mathcal{W}^*$ with $W_i \cap W_{i+1} \neq \emptyset$ ($i = 0, \dots, n-1$). Since $W_{\mathcal{C}}$ is connected, there is at least one chain from W to W' .

Lemma 4.17. *Λ is path connected.*

Proof. Let $W, W' \in \mathcal{W}^*$ with $W \cap \overline{C}_v, W' \cap \overline{C}_v \neq \emptyset$, and $W = W_0, W_1, \dots, W_n = W'$ a chain from W to W' with the smallest possible n . Choose $x_0 \in W \cap \overline{C}_v$ and $x_{n+1} \in W' \cap \overline{C}_v$ so that x_0 does not lie in any wall other than W and x_{n+1} does not lie in any wall other than W' . Pick $x_i \in W_{i-1} \cap W_i$ ($i = 1, \dots, n$) and let σ be the path defined by $\sigma = x_0 x_1 * \dots * x_n x_{n+1}$. Note the lemma follows if $\sigma \subset \overline{C}_v$.

We show $\sigma \subset \overline{C}_v$ by inducting on the length of a chain. We first show $x_0 x_1 \subset \overline{C}_v$. It suffices to show $x_1 \in \overline{C}_v$ since $\overline{C}_v$ is convex. Assume $x_1 \notin \overline{C}_v$. Then there is a wall $\tilde{W} \neq W, W_1$ so that x_0 and x_1 lie in different components of $X - \tilde{W}$. Since $x_0 x_1 \subset W$, $W \cap \tilde{W} \neq \emptyset$. $\tilde{W} \neq W'$ holds, otherwise by the choice of n we have $\tilde{W} = W_1$, a contradiction. It follows that x_0 and x_{n+1} lie on the same side of $\tilde{W}$. Thus the part of σ from x_1 to x_{n+1} must cross $\tilde{W}$. Suppose $\tilde{W} \cap x_i x_{i+1} \neq \emptyset$ for some i , $1 \leq i \leq n$. Since $x_i x_{i+1} \subset W_i$ we have $\tilde{W} \cap W_i \neq \emptyset$. If $i > 2$, then the sequence $W, \tilde{W}, W_i, \dots, W_n = W'$ is a chain from W to W' with length less than n , contradicting to the choice of n . If $i = 1$, then $W, W_1, \tilde{W}$ is a 3-cycle, contradicting to Proposition 2.9. Therefore $i = 2$ and $\tilde{W} \cap W_2 \neq \emptyset$. If $\tilde{W} = W_2$, then the sequence $W, \tilde{W}, W_3, \dots, W_n = W'$ is a chain from W to W'

with length $n - 1$, again contradicting to the choice of n . Thus $i = 2$ and $\tilde{W} \neq W_2$. But then $W, W_1, W_2, \tilde{W}$ is a 4-cycle, contradicting to Proposition 2.9.

Now $x_1 \in \overline{C}_v$ and $x_1 \in W_0 \cap W_1$ imply $W_1 \cap \overline{C}_v \neq \phi$. An argument similar to the one in the preceding paragraph shows x_2 and x_{n+1} lie on the same side of W . It follows that x_2 and C_v lie on the same side of W . Therefore an initial open segment of x_1x_2 lies in $\overline{C}_v$ and does not intersect any wall other than W_1 . Choose a point x'_1 belonging to this initial open segment of x_1x_2 . Now consider the sequence $W_1, W_2, \dots, W_n$ and let σ' be the part of σ from x'_1 to x_{n+1} . The induction hypothesis implies that $\sigma' \subset \overline{C}_v$. Now the lemma follows. $\square$

By Theorem 2.5 $\overline{C}_v / \text{Stab}_\Gamma \overline{C}_v$ is compact. Lemma 4.17 implies Λ is a boundary component of $\overline{C}_v$. It follows that $\Lambda / \text{Stab}_\Gamma \Lambda$ is a closed 3-manifold, being a quotient of a boundary component of $\overline{C}_v / \text{Stab}_\Gamma \overline{C}_v$.

Next we construct a graph G associated to Λ . The vertex set of G is in one-to-one correspondence with $\{W \cap \overline{C}_v : W \cap \overline{C}_v \neq \phi, W \in \mathcal{W}^*\}$. Let v_1 and v_2 be two vertices of G corresponding to $W_1 \cap \overline{C}_v$ and $W_2 \cap \overline{C}_v$ respectively. There is an edge connecting v_1 and v_2 if and only if $W_1 \cap W_2 \cap \overline{C}_v \neq \phi$.

The following lemma is clear from the definitions.

Lemma 4.18. *The graph G is a tree if there is no cycle in X .*

Lemma 4.19. *Λ is simply connected if there is no cycle in X .*

Proof. We notice G is the nerve of the covering $\{W \cap \overline{C}_v\}$ of Λ since $W_1 \cap W_2 \cap W_3 = \phi$ if W_1, W_2, W_3 are distinct. All $W \cap \overline{C}_v$ and their nonempty intersections are convex and thus contractible. Therefore G is homotopy equivalent to Λ . The lemma now follows from Lemma 4.18. $\square$

Suppose there is no cycle in X . Set $\Gamma_1 = \text{Stab}_\Gamma \Lambda$. Lemma 4.19 implies that $\Gamma_1 \cong \pi_1(\Lambda / \Gamma_1)$ and so $H \subset \Gamma_1$ is a subgroup of the fundamental group of a closed 3-manifold. We may assume Λ / Γ_1 is orientable by replacing Γ_1 with an index two subgroup if necessary. Λ / Γ_1 is a Haken manifold: Pick any $W_1 \neq W_2 \in \mathcal{W}^*$ with $W_1 \cap W_2 \neq \phi$ and $W_i \cap \overline{C}_v \neq \phi$ ($i = 1, 2$); then $F := W_1 \cap W_2$ is a 2-flat contained in Λ and $F / \text{Stab}_{\Gamma_1} F$ is a torus or Klein bottle embedded in Λ / Γ_1 ; the inclusion of $F / \text{Stab}_{\Gamma_1} F$ into Λ / Γ_1 clearly induces an injective homomorphism on the fundamental groups. Now Theorem 3.3 implies H has the TA-property since any solvable subgroup of a group acting properly and cocompactly on a Hadamard space is virtually free abelian. The proof of Theorem 4.1 is now complete.

5. CYCLES OF HIGHER RANK SUBMANIFOLDS

Throughout this section let $M = X / \Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature, and $H \subset \Gamma$ a subgroup as described at the beginning of Section 4.3. Recall H contains no Type D isometries. We proved in Section 4.3 that H has the TA-property if there is no cycle in X . In this section we discuss the Tits alternative without assuming the nonexistence of cycles.

Recall a singular geodesic in X is a geodesic of the form $\{q\} \times R \subset Q \times R = W$, where $q \in Q$ and W is a wall. When W_1 and W_2 are two walls and $F = W_1 \cap W_2$ is a 2-flat, there are two families of parallel singular geodesics in F . The angle α_F ($0 < \alpha_F \leq \frac{\pi}{2}$) between them is a *singular angle*. Since modulo Γ there are only a finite number of walls in X , there are only a finite number of singular angles.

Now we are ready to state the main result of this section.

Theorem 5.1. *Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature. Suppose there is a number α , $\frac{2\pi}{5} < \alpha \leq \frac{\pi}{2}$ so that all the singular angles of X are equal to α . If a subgroup of the fundamental group of M is not virtually free abelian, then it contains a free group of rank two.*

Remark 5.2. U. Abresch and V. Schroeder ([AS]) constructed a class of real analytic 4-manifolds of nonpositive sectional curvature, where all the singular angles are $\frac{\pi}{2}$.

5.1. Incidence Graph. We use the notation of Section 4.3. We first construct a graph G_* which reflects the incidence relation of the members of $\mathcal{W}^*$. The vertex set of G_* is in one-to-one correspondence with $\mathcal{W}^*$. Two vertices are joined by an edge if the corresponding walls have nonempty intersection. G_* is a connected graph. The vertex corresponding to the wall W is still denoted by W , and the edge joining two vertices W_1, W_2 is denoted by W_1W_2 . We declare each edge of G_* has length 1 and let d_* be the induced path metric on G_* . Since by Proposition 2.9 there are no n -cycles in X for $n \leq 4$, we have:

Lemma 5.3. *Any injective loop in G_* has length at least 5.*

The lemma in particular implies for $W_1, W_2 \in \mathcal{W}^*$ with $d_*(W_1, W_2) = 2$, there is a unique $W \in \mathcal{W}^*$ with $d_*(W_1, W) = d_*(W, W_2) = 1$.

Note $h(\mathcal{W}^*) = \mathcal{W}^*$ for each $h \in H$. Hence H induces an action on G_* . For $h \in H$ we denote by h_* the isomorphism of G_* induced by h .

Lemma 5.4. *If H has a fixed point in G_* , then H has the TA-property.*

Proof. Since G_* is a graph, after passing to an index two subgroup if necessary, we may assume H fixes a vertex W of G_* . Then $h(W) = W$ for all $h \in H$, here $W \subset X$ denotes the wall in X . Now the lemma follows from the proof of Lemma 4.15. □

5.2. Tits Geodesics. We continue to use the notation of Section 4.3. In this section we take a close look at Tits geodesics in $\mathcal{C} = \cup_{W \in \mathcal{W}^*} \partial_T W$.

Let W be a wall. A singular geodesic $c = \{q\} \times R \subset Q \times R = W$ determines two points $c(+\infty)$, $c(-\infty)$ in $\partial_\infty W$. Set $w(+\infty) = c(+\infty)$, $w(-\infty) = c(-\infty)$ and call $w(+\infty)$, $w(-\infty)$ the *poles* of W .

The following proposition follows from the results in [HS1].

Proposition 5.5. *Let $W_1, W_2 \in \mathcal{W}^*$ be two walls.*

- (i) *If $d_*(W_1, W_2) = 1$, i.e., if $W_1 \cap W_2 = F$ is a 2-flat, then $\partial_\infty W_1 \cap \partial_\infty W_2 = \partial_\infty F$;*
- (ii) *If $d_*(W_1, W_2) = 2$, then $\partial_\infty W_1 \cap \partial_\infty W_2 = \{w(+\infty), w(-\infty)\}$ where W is the unique wall with $d_*(W_1, W) = d_*(W, W_2) = 1$;*
- (iii) *If $d_*(W_1, W_2) \geq 3$, then $\partial_\infty W_1 \cap \partial_\infty W_2 = \emptyset$.*

Let $W_1, W_2 \in \mathcal{W}^*$ with $d_*(W_1, W_2) = 1$. Then $W_1 \cap W_2 = F$ is a 2-flat and $\partial_T W_1 \cap \partial_T W_2 = \partial_T F$ is isometric to the unit circle. $\partial_T F$ admits a unique metric graph structure with vertex set $\{w_1(+\infty), w_1(-\infty), w_2(+\infty), w_2(-\infty)\}$.

Let $W = Q \times R \in \mathcal{W}^*$. Since W is closed and Q is a nonflat Hadamard 2-manifold, $\partial_T W$ admits a unique metric graph structure with the following properties:

- (a) the vertex set is $\{w'(+\infty), w'(-\infty) : d_*(W', W) \leq 1\}$;
- (b) for each wall W' with $d_*(W', W) = 1$, the inclusion $\partial_T(W' \cap W) \subset \partial_T W$ is an isometric embedding between metric graphs;
- (c) each edge connecting the two poles $w(+\infty)$, $w(-\infty)$ has length π .

It follows from the results in [HS1] that $\mathcal{C}$ admits a unique metric graph structure with vertex set $\{w(+\infty), w(-\infty) : W \in \mathcal{W}^*\}$ such that for each $W \in \mathcal{W}^*$, the inclusion $\partial_T W \subset \mathcal{C}$ is an isometric embedding between metric graphs.

We next look at how Tits geodesics in $\mathcal{C}$ travel between different $\partial_\infty W$, $W \in \mathcal{W}^*$.

Lemma 5.6. *Let $W \in \mathcal{W}^*$, $\sigma : [a, b] \rightarrow \partial_T X$ a minimal geodesic and $t_0 \in (a, b)$. If there is some $\epsilon > 0$ such that $\sigma(t) \in \partial_\infty W$ for $t \in (t_0 - \epsilon, t_0]$ and $\sigma(t) \notin \partial_\infty W$ for $t \in (t_0, t_0 + \epsilon)$, then $\sigma(t) \notin \partial_\infty W$ for $t \in (t_0, b]$.*

Proof. Suppose the lemma is false and let $t_1 = \min\{t \in (t_0, b] : \sigma(t) \in \partial_\infty W\}$. Notice the minimal in the definition of t_1 makes sense since $\partial_\infty W$ is closed in $\partial_\infty X$. By definition $\partial_\infty W \cap \sigma_{|[t_0, t_1]} = \emptyset$. Since $\sigma(t_0), \sigma(t_1) \in \partial_\infty W$ we have $d_T(\sigma(t_0), \sigma(t_1)) \leq \pi$. Let $\sigma' \subset \partial_T W$ be a minimal geodesic from $\sigma(t_0)$ to $\sigma(t_1)$. Then $\sigma' \cup \sigma_{|[t_0, t_1]}$ is a closed geodesic in the CAT(1) space $\partial_T X$ with length

$$\text{length}(\sigma') + \text{length}(\sigma_{|[t_0, t_1]}) = 2d_T(\sigma(t_0), \sigma(t_1)) \leq 2\pi.$$

It follows that $d_T(\sigma(t_0), \sigma(t_1)) = \pi$. Since $\sigma_{|[t_0 - \epsilon, t_0]} \subset \partial_\infty W$, the description of $\partial_T W$ shows $d_T(\sigma(t), \sigma(t_1)) < \pi$ for $t \in (t_0 - \epsilon, t_0)$, contradicting to the fact that σ is minimal. $\square$

The following proposition follows from Lemma 5.6 and the description of $\mathcal{C}$.

Proposition 5.7. *Given any minimal geodesic $\sigma : [a, b] \rightarrow \mathcal{C}$, there are walls $W_1, W_2, \dots, W_n$ and numbers $a < t_1 < t_2 < \dots < t_n < b$ with the following properties:*

- (i) $\{\sigma(t_i), 1 \leq i \leq n\}$ is the set of poles in the interior of σ ;
- (ii) $\sigma(t_i)$ is a pole of W_i for each $1 \leq i \leq n$;
- (iii) the sequence of walls $W_1, W_2, \dots, W_n$ determines an injective edge path in G_* .

5.3. Subgroups Containing Type C Isometries. In this section we show the subgroup $H \subset \Gamma$ has the TA-property if the singular angles are large and H contains a Type C isometry.

We recall there are only a finite number of singular angles.

Proposition 5.8. *Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature. Suppose all the singular angles of X are strictly larger than $\frac{\pi}{3}$. If $H \subset \Gamma$ is a subgroup as described at the beginning of Section 4.3 and contains a Type C isometry, then H has the TA-property.*

Proof. For any two Type C isometries $h, \tilde{h} \in H$, we set $n(h, \tilde{h}) = d_*(W, \tilde{W})$ where $W = P(h)$ and $\tilde{W} = P(\tilde{h})$. Notice $h(+\infty), h(-\infty)$ are the poles of W , and $\tilde{h}(+\infty), \tilde{h}(-\infty)$ are the poles of $\tilde{W}$. First suppose there are two Type C isometries h and $\tilde{h}$ with $n(h, \tilde{h}) \geq 3$. For any $\xi \in \{h(+\infty), h(-\infty)\}$ and $\eta \in \{\tilde{h}(+\infty), \tilde{h}(-\infty)\}$, let $\sigma : [0, a] \rightarrow \partial_T X$ be a minimal geodesic from ξ to η . If there is no pole in the interior of σ , then $\sigma \subset \partial_\infty W \cap \partial_\infty \tilde{W}$, contradicting to Proposition 5.5. Let $\sigma(t_i)$ ($i = 1, 2, \dots, n$) with $0 < t_1 < t_2 < \dots < t_n < a$ be all the poles in the interior of σ . If $n = 1$, then $\sigma(t_1) \in \partial_\infty W \cap \partial_\infty \tilde{W}$, again contradicting to Proposition 5.5. Therefore there are at least two poles in the interior of σ . Since the endpoints of σ are also poles, σ contains at least 4 poles. By assumption on the singular angles, the Tits distance between any two distinct poles is $> \frac{\pi}{3}$. It follows that $d_T(\xi, \eta) > \pi$. By Theorem 2.14 $\langle h, \tilde{h} \rangle$ contains a free group of rank two.

Now suppose there are two Type C isometries h and $\tilde{h}$ with $n(h, \tilde{h}) = 2$. Let $W_1 = P(h)$, $W_2 = P(\tilde{h})$. There exists an integer $k \geq 1$ with $\text{Min}(h^k) = W_1$ and $\text{Min}(\tilde{h}^k) = W_2$. By Lemma 5.3 there is a unique wall W with $W \cap W_1 \neq \emptyset$ and $W \cap W_2 \neq \emptyset$. We have $h^k(W) = W$, $\tilde{h}^k(W) = W$. Let $W \cap W_1 = c_1 \times R \subset Q \times R = W$ and $W \cap W_2 = c_2 \times R \subset Q \times R = W$, where c_1, c_2 are complete geodesics in Q . Theorem 2.5 implies $\{c_1(+\infty), c_1(-\infty)\} \cap \{c_2(+\infty), c_2(-\infty)\} = \emptyset$. h^k acts on W as $h^k = (h_1, t) : Q \times R \rightarrow Q \times R$, where $h_1 : Q \rightarrow Q$ is a hyperbolic isometry of Q with c_1 as an axis and t is a translation of R . Similarly $\tilde{h}^k$ acts on W as $\tilde{h}^k = (h_2, t')$ where h_2 is a hyperbolic isometry of Q with c_2 as an axis and t' is a translation of R . Since Q is nonflat and admits a cocompact group of isometries, Q is hyperbolic in the sense of Gromov. Theorem 2.11 implies $\langle h_1, h_2 \rangle$ contains a free group of rank two. It follows that $\langle h^k, \tilde{h}^k \rangle$ contains a free group of rank two.

Now suppose $n(h, \tilde{h}) \leq 1$ for any two Type C isometries $h, \tilde{h} \in H$. For any three Type C isometries $h_1, h_2, h_3 \in H$, let $W_i = P(h_i)$ ($i = 1, 2, 3$). If $W_1 \neq W_2$ and $W_1 \neq W_3$, then $W_2 =$

W_3 since $n(h_2, h_3) \leq 1$ and G_* has no injective edge loop with length 3. It follows that the set $\{P(h) : h \in H \text{ is of Type C}\}$ consists of one wall or two intersecting walls. Notice the set $\{P(h) : h \in H \text{ is of Type C}\}$ is invariant under the action of H since the conjugate of a Type C isometry is still a Type C isometry and $P(ghg^{-1}) = g(P(h))$. The proposition now follows from Lemma 5.4. $\square$

5.4. Admissible Subsets. For a Hadamard space X , $\xi \in \partial_T X$ and $r > 0$, we let $\overline{B}(\xi, r)$ be the closed metric ball with center ξ and radius r : $\overline{B}(\xi, r) = \{\eta \in \partial_T X : d_T(\xi, \eta) \leq r\}$.

Definition 5.9. Let X be a Hadamard space and $H \subset \text{Isom}(X)$. A nonempty H -invariant subset $M \subset \partial_\infty X$ is H -admissible if $M \subset \overline{B}(h(+\infty), \pi) \cap \overline{B}(h(-\infty), \pi)$ for each hyperbolic isometry $h \in H$.

Proposition 5.10. Let $M = X/\Gamma$ be a rank 1 closed real analytic 4-manifold of nonpositive sectional curvature, and $H \subset \Gamma$ a subgroup as described at the beginning of Section 4.3. Then there exists a H -admissible subset.

Proof. We notice $h(+\infty), h(-\infty) \in \Lambda(H)$ for any $h \in H$. Let $M \subset \Lambda(H)$ be a H -minimal set. Since by assumption H does not contain any rank one isometries, Proposition 2.18 implies M is H -admissible. $\square$

5.5. Intersection of Tits Balls. In order to study H -admissible subsets in $\partial_\infty X$, we need to look at the intersection of $\overline{B}(h(+\infty), \pi)$ and $\overline{B}(h(-\infty), \pi)$ for $h \in H$.

Call $h \in H$ a *squared Type B isometry* if $h = g^2$ for a Type B isometry $g \in H$. Clearly $h \in H$ is a squared Type B isometry if and only if any of its conjugates is a squared Type B isometry. Notice for a squared Type B isometry h , $\text{Min}(h) = P(h)$. If $h \in H$ is of Type A, then $\text{Fix}(h_*)$ is the vertex W in G_* , where W is the only wall containing $\text{Min}(h) \subset X$. If $h \in H$ is a squared Type B isometry, then $\text{Fix}(h_*)$ is the edge $W_1 W_2$ in G_* , where W_1, W_2 are the two walls with $W_1 \cap W_2 = \text{Min}(h) \subset X$. For any $h \in H$ and any wall W , set

$$\overline{B}(h, W) = \overline{B}(h(+\infty), \pi) \cap \overline{B}(h(-\infty), \pi) \cap \partial_\infty W.$$

Lemma 5.11. Suppose the singular angles of X are all equal to $\alpha > \frac{2}{5}\pi$. Let $h \in H$ be a Type A or squared Type B isometry, and $W \in \mathcal{W}^*$ with $d_*(W, \text{Fix}(h_*)) \geq 3$. Then $\overline{B}(h, W) = \phi$.

Proof. Suppose the lemma is false and pick $\xi \in \overline{B}(h, W)$. Let $c : [0, a] \rightarrow \partial_T X$ ($a \leq \pi$) be a minimal geodesic from $h(+\infty)$ to ξ , $W_1, \dots, W_k$ a sequence of walls and $\{t_i\}$ ($0 < t_1 < \dots < t_k < a$) a sequence of numbers as in Proposition 5.7. By definition of Type A and squared Type B isometries, $h(+\infty)$ is not a pole and $h_*(W_1) = W_1$. $a \leq \pi$ implies $k \leq 3$ since by assumption the Tits distance between any two distinct poles is at least $\frac{2}{5}\pi$. If there is no pole in the interior of c , then there is a wall $\tilde{W}$ with $h_*(\tilde{W}) = \tilde{W}$ and $d_*(\tilde{W}, W) \leq 1$, contradicting to the assumption $d_*(W, \text{Fix}(h_*)) \geq 3$. If $k = 1$, then $\xi \in \partial_\infty W_1 \cap \partial_\infty W$ and by Proposition 5.5 we have $d_*(W_1, W) \leq 2$, again contradicting to the assumption. If $k = 2$, then $\xi \in \partial_\infty W_2 \cap \partial_\infty W$ and since $d_*(W, \text{Fix}(h_*)) \geq 3$ we have $d_*(W_2, W) = 2$ and ξ must be a pole. In this case let W_3 be the wall with ξ as one of its two poles and set $t_3 = a$. If $k = 3$, then ξ can not be a pole since $a \leq \pi$ and the Tits distance between any two poles is at least $\frac{2}{5}\pi$. In any case we have $\xi \in B(c(t_3), \frac{\pi}{5})$. Similarly if $c' : [0, b] \rightarrow \partial_T X$ ($b \leq \pi$) is a minimal geodesic from $h(-\infty)$ to ξ , we have walls W'_1, W'_2, W'_3 ($h_*(W'_1) = W'_1$) and poles $c'(t'_1), c'(t'_2), c'(t'_3)$ such that $\xi \in B(c'(t'_3), \frac{\pi}{5})$. It follows from the triangle inequality that $d_T(c(t_3), c'(t'_3)) < \frac{2}{5}\pi$. The assumption on singular angles implies $c(t_3) = c'(t'_3)$. Consequently $W'_3 = W_3$.

We next show $W'_1 = W_1$. Note $t_1, t'_1 < \frac{\pi}{5}$. Since $d_T(h(+\infty), h(-\infty)) = \pi$, triangle inequality implies $d_T(c(t_1), c'(t'_1)) > \frac{3}{5}\pi$. The two poles $c(t_1), c'(t'_1)$ both lie on $\partial_\infty W_1$. Notice for any pole η in

$\partial_\infty W_1$, we have $d_T(\eta, c(t_1)) \in \{\alpha, \pi - \alpha, \pi\}$. The assumption on α now implies $d_T(c(t_1), c'(t'_1)) = \pi$. Therefore $c'(t'_1)$ is also a pole of W_1 , and $W'_1 = W_1$.

Now we have two injective edge paths of length 2 from $W_1 = W'_1$ to $W_3 = W'_3$: $W_1 W_2 W_3$, $W'_1 W'_2 W'_3$. By Lemma 5.3 we have $W_2 = W'_2$. Notice $t_3 - t_2 = \alpha$ otherwise $t_3 - t_2 \geq \pi - \alpha$ and $a > (t_3 - t_2) + (t_2 - t_1) \geq (\pi - \alpha) + \alpha = \pi$. The same argument also shows $\alpha < \frac{\pi}{2}$ and $t'_3 - t'_2 = \alpha$. Now the three poles $c(t_2)$, $c'(t'_2)$, $c(t_3)$ all lie on the circle $\partial_\infty W_2 \cap \partial_\infty W_3$ and $d_T(c(t_2), c(t_3)) = d_T(c'(t'_2), c(t_3)) = \alpha$. Since $\alpha < \frac{\pi}{2}$ we have $c'(t'_2) = c(t_2)$. Similarly we conclude $c'(t'_1) = c(t_1)$.

Since $t_1, t'_1 < \frac{\pi}{5}$, by triangle inequality

$$d_T(h(+\infty), h(-\infty)) \leq d_T(h(+\infty), c(t_1)) + d_T(c'(t'_1), h(-\infty)) < \frac{2}{5}\pi,$$

contradicting to the fact $d_T(h(+\infty), h(-\infty)) = \pi$. □

If $h \in H$ is of Type A and W is a wall with $d_*(W, \text{Fix}(h_*)) = 2$, there is exactly one wall W' with $d_*(W, W') = d_*(W', \text{Fix}(h_*)) = 1$. If $h \in H$ is a squared Type B isometry and W is a wall with $d_*(W, \text{Fix}(h_*)) = 2$, then the set

$$\{W' : d_*(W, W') = d_*(W', \text{Fix}(h_*)) = 1\}$$

consists of either one or two elements. These assertions follow from Lemma 5.3.

Lemma 5.12. *Let $h \in H$ be a Type A or squared Type B isometry, and $W \in \mathcal{W}^*$ such that the following holds: $d_*(W, \text{Fix}(h_*)) = 2$. Suppose the singular angles of X are all equal to $\alpha > \frac{2}{5}\pi$.*

(i) *If there is only one wall W' with $d_*(W, W') = d_*(W', \text{Fix}(h_*)) = 1$, then*

$$\overline{B}(h, W) \subset \partial_\infty W \cap \partial_\infty W' - \{w(+\infty), w(-\infty)\};$$

(ii) *If there are two walls W_1, W_2 with $d_*(W, W_i) = d_*(W_i, \text{Fix}(h_*)) = 1$, then*

$$\overline{B}(h, W) \subset (\partial_\infty W \cap \partial_\infty W_1) \cup (\partial_\infty W \cap \partial_\infty W_2) - \{w(+\infty), w(-\infty)\}.$$

Proof. We fix an arbitrary $\xi \in \overline{B}(h, W)$. Let $c : [0, a] \rightarrow \partial_T X$ be a minimal geodesic from $h(+\infty)$ to ξ , and $c' : [0, b] \rightarrow \partial_T X$ a minimal geodesic from $h(-\infty)$ to ξ , where $a, b \leq \pi$. Let $W_1, \dots, W_k$ be a sequence of walls and $\{t_i\}$ ($0 < t_1 < \dots < t_k < a$) a sequence of numbers provided by Proposition 5.7 corresponding to c , and $W'_1, \dots, W'_{k'}$ and $\{t'_i\}$ ($0 < t'_1 < \dots < t'_{k'} < b$) corresponding to c' . As in the proof of Lemma 5.11 we see $h_*(W_1) = W_1$, $h_*(W'_1) = W'_1$ and $1 \leq k \leq 3$, $1 \leq k' \leq 3$.

If $k = 1$, then $\xi \in \partial_\infty W_1 \cap \partial_\infty W$. Since $d_*(W, \text{Fix}(h_*)) = 2$ and $h_*(W_1) = W_1$, Proposition 5.5 implies ξ is a pole of W' , where W' is the only wall with $d_*(W, W') = 1$ and $d_*(W', W_1) = 1$. We clearly have $\xi \in \partial_\infty W \cap \partial_\infty W' - \{w(+\infty), w(-\infty)\}$ in this case. From now on we assume $2 \leq k, k' \leq 3$.

Suppose $k = 2$. In this case $\xi \in \partial_\infty W_2 \cap \partial_\infty W$. If ξ is not a pole, then W_2 is the only wall with $d_*(W, W_2) = d_*(W_2, W_1) = 1$ and clearly

$$\xi \in \partial_\infty W \cap \partial_\infty W_2 - \{w(+\infty), w(-\infty)\}.$$

Suppose ξ is a pole of some wall W_3 . Then the assumption on α implies $k' = 2$. In this case we have two injective edge paths in G_* : $W_1 W_2 W_3$ and $W'_1 W'_2 W'_3$. The proof of Lemma 5.11 yields a contradiction if $W'_1 = W_1$. Hence $W'_1 \neq W_1$, h is a Type B isometry and $\text{Fix}(h_*) = W_1 W'_1$. Notice $t_1 < \frac{\pi}{5}$ since $a \leq \pi$ and $c(t_1), c(t_2), \xi$ are three poles on c . Similarly $t'_1 < \frac{\pi}{5}$. Recall $c(t_1)$ is a pole of W_1 . Let $p \neq c(t_1)$ be the other pole of W_1 . Since $h(+\infty), h(-\infty), p$ and $c(t_1)$ all lie on the circle $\partial_T W_1 \cap \partial_T W'_1$, we see $d_T(h(-\infty), p) = d_T(h(+\infty), c(t_1)) = t_1$. It follows that

$$d_T(c'(t'_1), p) \leq d_T(c'(t'_1), h(-\infty)) + d_T(h(-\infty), p) = t'_1 + t_1 < \frac{2}{5}\pi.$$

As $c'(t'_1)$ and p are poles of two distinct walls W'_1 and W_1 respectively, we have $c'(t'_1) \neq p$ and $d_T(c'(t'_1), p) \geq \alpha > \frac{2}{5}\pi$, a contradiction.

Now we assume $k = k' = 3$. In this case ξ is not a pole. The proof of Lemma 5.11 shows $c(t_3) = c'(t'_3)$ and $W_3 = W'_3$. If $W'_1 = W_1$ then the proof of Lemma 5.11 yields a contradiction. So $W'_1 \neq W_1$, $Fix(h_*) = W_1 W'_1$ and h is a Type B isometry. Now consider $c(t_3) = c'(t'_3) \in \overline{B}(h, W_3)$ instead of ξ and the preceding paragraph yields a contradiction. $\square$

5.6. Proof of Theorem 5.1. In this section we finish the proof of Theorem 5.1. By Lemma 5.4 and Proposition 5.8, we may assume H does not have a global fixed point in G_* and each nontrivial $h \in H$ is of Type A or Type B.

Let $M \subset \partial_\infty X$ be a H -admissible subset. By Lemma 5.11, if W is a wall with $d_*(W, Fix(h_*)) \geq 3$ for some $h \in H$, then $M \cap \partial_\infty W = \phi$.

Lemma 5.13. *Let $M \subset \partial_\infty X$ be H -admissible. Suppose W is a wall such that $Fix(h_*) = \{W\}$ for a Type A isometry $h \in H$. Then $M \cap \partial_\infty W = \phi$.*

Proof. Since $Fix((ghg^{-1})_*) = g_*(Fix(h_*))$ and the conjugate of a Type A isometry is still a Type A isometry, the assumption that H does not have a global fixed point in G_* implies that there is a wall $W' \neq W$ with $Fix(h'_*) = \{W'\}$ for a Type A isometry $h' \in H$. If $d_*(W', W) \geq 3$, then Lemma 5.11 implies $\overline{B}(h', W) = \phi$. By the definition of a H -admissible set, $M \subset \overline{B}(h'(+\infty), \pi) \cap \overline{B}(h'(-\infty), \pi)$ and thus $M \cap \partial_\infty W \subset \overline{B}(h'(+\infty), \pi) \cap \overline{B}(h'(-\infty), \pi) \cap \partial_\infty W = \overline{B}(h', W) = \phi$. From now on we assume $d_*(W', W) \leq 2$.

Suppose $d_*(W', W) = 1$. Then $h'(W) \neq W$ and $d_*(W, W') = d_*(h'(W), W') = 1$. By Lemma 5.3 $d_*(W, h'(W)) = 2$. Thus we may assume $d_*(W', W) = 2$. Then $d_*(h(W'), W) = 2$. Let W_1, W_2 be the unique walls with $d_*(W, W_1) = d_*(W_1, W') = 1$, $d_*(W, W_2) = d_*(W_2, h(W')) = 1$. Notice $W_2 = h(W_1) \neq W_1$. Lemma 5.12 implies

$$\overline{B}(h', W) \subset \partial_\infty W \cap \partial_\infty W_1 - \{w(+\infty), w(-\infty)\}$$

and

$$\overline{B}(hh'h^{-1}, W) \subset \partial_\infty W \cap \partial_\infty W_2 - \{w(+\infty), w(-\infty)\}.$$

M is H -admissible implies

$$\begin{aligned} M \cap \partial_\infty W &\subset \overline{B}(h', W) \cap \overline{B}(hh'h^{-1}, W) \subset \\ &\subset (\partial_\infty W \cap \partial_\infty W_1 - \{w(+\infty), w(-\infty)\}) \cap (\partial_\infty W \cap \partial_\infty W_2 - \{w(+\infty), w(-\infty)\}) \subset \\ &\subset \partial_\infty W_1 \cap \partial_\infty W_2 - \{w(+\infty), w(-\infty)\} = \phi. \end{aligned}$$

The last equality follows from Proposition 5.5 and the facts that $W_1 \neq W_2$ and $d_*(W_1, W) = 1$, $d_*(W, W_2) = 1$. $\square$

Lemma 5.14. *Let $M \subset \partial_\infty X$ be H -admissible. Suppose W is a wall with $d_*(Fix(h_*), W) = 1$ for a Type A isometry $h \in H$. Then $M \cap \partial_\infty W = \phi$.*

Proof. Suppose $M \cap \partial_\infty W \neq \phi$. Let $Fix(h_*) = \{W_0\}$. The argument in the proof of Lemma 5.13 shows that there is a wall W' and a Type A isometry $h' \in H$ such that $Fix(h'_*) = \{W'\}$ and $d_*(W_0, W') = 2$ or 3 .

Let us first assume $d_*(W_0, W') = 3$. Then $M \cap \partial_\infty W \neq \phi$ implies $d_*(W, W') = 2$ and $d_*(W, h(W')) = 2$. Let W_1 be the unique wall with $d_*(W, W_1) = d_*(W_1, W') = 1$, and W_2 the unique wall with $d_*(W, W_2) = d_*(W_2, h(W')) = 1$. Lemma 5.12 implies

$$\begin{aligned} \phi &\neq M \cap \partial_\infty W \subset \overline{B}(h', W) \cap \overline{B}(hh'h^{-1}, W) \subset \\ &\subset (\partial_\infty W \cap \partial_\infty W_1 - \{w(+\infty), w(-\infty)\}) \cap (\partial_\infty W \cap \partial_\infty W_2 - \{w(+\infty), w(-\infty)\}) \subset \\ &\subset \partial_\infty W_1 \cap \partial_\infty W_2 - \{w(+\infty), w(-\infty)\}, \end{aligned}$$

and therefore $W_1 = W_2$. Since M is H -invariant, $M \cap \partial_\infty W \neq \phi$ implies $M \cap \partial_\infty h(W)$ is nonempty. The above argument applied to $h(W)$ instead of W shows there exists a wall W'_1 with $d_*(h(W), W'_1) =$

$d_*(W'_1, W') = d_*(W'_1, h(W')) = 1$. If $W'_1 \neq W_1$, then $W_1 W' W'_1 h(W') W_1$ is an injective edge loop with length 4, a contradiction. If $W'_1 = W_1$, then $W_0 W W_1 h(W) W_0$ is an injective edge loop with length 4, again a contradiction.

Now assume $d_*(W_0, W') = 2$. $M \cap \partial_\infty W \neq \emptyset$, Lemma 5.13 and the remark before Lemma 5.13 imply $d_*(W, W') = 1$ or 2. By replacing W' with $h(W')$ if necessary we may assume $d_*(W, W') = 2$. The argument in the preceding paragraph again yields a contradiction. $\square$

Lemma 5.15. *Let $M \subset \partial_\infty X$ be H -admissible. Suppose W is a wall with $d_*(\text{Fix}(h_*), W) = 2$ for a Type A isometry $h \in H$. Then $M \cap \partial_\infty W = \emptyset$.*

Proof. Let $\text{Fix}(h_*) = \{W_0\}$ and W_1 be the unique wall with $d_*(W_0, W_1) = 1$ and $d_*(W_1, W) = 1$. Then by Lemma 5.12

$$M \cap \partial_\infty W \subset \overline{B}(h, W) \subset \partial_\infty W_1 \cap \partial_\infty W - \{w(+\infty), w(-\infty)\} \subset \partial_\infty W_1.$$

But Lemma 5.14 implies $M \cap \partial_\infty W_1 = \emptyset$. It follows $M \cap \partial_\infty W \subset M \cap \partial_\infty W_1 = \emptyset$. $\square$

Lemmas 5.13, 5.14, 5.15, Proposition 5.10 and the remark before Lemma 5.13 together imply that H contains no Type A isometries. From now on we assume each nontrivial $h \in H$ is of Type B and H does not have a global fixed point in G_* . Recall $h \in H$ is a squared Type B isometry if and only if any of its conjugates is a squared Type B isometry.

Lemma 5.16. *If there are squared Type B isometries $h, h' \in H$ with $d_*(\text{Fix}(h_*), \text{Fix}(h'_*)) \geq 3$, then H contains a free group of rank two.*

Proof. Let $\xi \in \{h(+\infty), h(-\infty)\}$, $\eta \in \{h'(+\infty), h'(-\infty)\}$ and $\sigma : [0, a] \rightarrow \partial_T X$ be a minimal geodesic from ξ to η . Let the walls $W_1, \dots, W_k$ and the numbers $0 < t_1 < \dots < t_k < a$ be provided by Proposition 5.7. Then $h_*(W_1) = W_1$ and $h'_*(W_k) = W_k$. Since $d_*(\text{Fix}(h_*), \text{Fix}(h'_*)) \geq 3$, we have $k \geq 4$. Since each singular angle equals $\alpha > \frac{2}{5}\pi$, the length of σ is at least $3 \times \frac{2}{5}\pi > \pi$. So $d_T(\xi, \eta) > \pi$ and the lemma now follows from Theorem 2.14. $\square$

Lemma 5.17. *Suppose each nontrivial $h \in H$ is of Type B and H does not have a global fixed point in G_* . Then there are squared Type B isometries $h, h' \in H$ with $d_*(\text{Fix}(h_*), \text{Fix}(h'_*)) \geq 2$.*

Proof. First suppose $d_*(\text{Fix}(h_*), \text{Fix}(h'_*)) = 0$ for all squared Type B isometries $h, h' \in H$. Fix a squared Type B isometry $h \in H$ and let $\text{Fix}(h_*) = W_1 W_2$. Since H does not fix the midpoint of $W_1 W_2$, there is some element g in H such that $g_*(W_1 W_2) \cap W_1 W_2 = \{W_1\}$ or $\{W_2\}$. After possibly relabeling W_1 and W_2 we may assume $g_*(W_1 W_2) \cap W_1 W_2 = \{W_1\}$. Set $h_1 = h$ and $h_2 = ghg^{-1}$. h_1 and h_2 are two squared Type B isometries with $\text{Fix}(h_{1*}) \cap \text{Fix}(h_{2*}) = \{W_1\}$. Since H does not fix W_1 , there is some $k \in H$ such that $k_*(W_1) \neq W_1$. Set $k_1 = kh_1k^{-1}$, $k_2 = kh_2k^{-1}$. Then k_1 and k_2 are two squared Type B isometries with $\text{Fix}(k_{1*}) \cap \text{Fix}(k_{2*}) = \{k_*(W_1)\}$. Now it is not hard to derive from Lemma 5.3 that there are $i, j \in \{1, 2\}$ such that $\text{Fix}(h_{i*}) \cap \text{Fix}(k_{j*}) = \emptyset$, a contradiction.

Now assume $d_*(\text{Fix}(h_*), \text{Fix}(h'_*)) \leq 1$ for any two squared Type B isometries $h, h' \in H$. By the above paragraph, there are two squared Type B isometries $g, h \in H$ with $d_*(\text{Fix}(h_*), \text{Fix}(g_*)) = 1$. Let $\text{Fix}(h_*) = W_2 W_3$, $g_*(W_1) = W_1$ with $d_*(W_1, W_2) = 1$. Notice $\text{Fix}((ghg^{-1})_*) = g(W_2)g(W_3)$. It follows from $d_*(\text{Fix}(h_*), \text{Fix}((ghg^{-1})_*)) \leq 1$ and Lemma 5.3 that $d_*(W_3, g(W_3)) = 1$. Similarly we have $d_*(W_3, g^2(W_3)) = 1$ and $d_*(g^2(W_3), g(W_3)) = 1$. It follows that we have an injective edge loop $W_3 g(W_3) g^2(W_3) W_3$ with length 3, a contradiction. $\square$

Lemma 5.18. *Let $M \subset \partial_\infty X$ be H -admissible, and $h, g \in H$ be two squared Type B isometries with $d_*(\text{Fix}(h_*), \text{Fix}(g_*)) = 2$ and $\text{Fix}(h_*) = WW'$. Then $M \cap \partial_\infty W = M \cap \partial_\infty W' = \phi$.*

Proof. We shall prove $M \cap \partial_\infty W = \phi$. The proof of $M \cap \partial_\infty W' = \phi$ is similar. If $d_*(W, \text{Fix}(g_*)) = 3$, then by Lemma 5.11 $M \cap \partial_\infty W \subset \overline{B}(g, W) = \phi$. Suppose $d_*(W, \text{Fix}(g_*)) = 2$. Then Lemma 5.12 implies that there is a set of walls $\{W_i\}_{i=1}^N$ ($N = 1$ or 2) with $d_*(W, W_i) = d_*(W_i, \text{Fix}(g_*)) = 1$ and

$$M \cap \partial_\infty W \subset \overline{B}(g, W) \subset \cup_{i=1}^N (\partial_\infty W \cap \partial_\infty W_i) - \{w(+\infty), w(-\infty)\}.$$

Fix an integer j with $\{W_1, W_N\} \cap \{h^j(W_1), h^j(W_N)\} = \phi$. Since $M \cap \partial_\infty W$ is also contained in $\overline{B}(h^j g h^{-j}, W) \subset \cup_{i=1}^N (\partial_\infty W \cap \partial_\infty h^j(W_i)) - \{w(+\infty), w(-\infty)\}$, it follows $M \cap \partial_\infty W$ is contained in the intersection of $\cup_{i=1}^N (\partial_\infty W \cap \partial_\infty W_i) - \{w(+\infty), w(-\infty)\}$ and $\cup_{i=1}^N (\partial_\infty W \cap \partial_\infty h^j(W_i)) - \{w(+\infty), w(-\infty)\}$, which is empty. $\square$

Lemma 5.19. *Let $M \subset \partial_\infty X$ be a H -admissible subset, $h, g \in H$ squared Type B isometries with $d_*(\text{Fix}(h_*), \text{Fix}(g_*)) = 2$, and W a wall with $d_*(W, \text{Fix}(h_*)) = 1$. Then $M \cap \partial_\infty W = \phi$.*

Proof. Notice $\text{Fix}((hgh^{-1})_*) = h_*(\text{Fix}(g_*))$ and by replacing h with h^2 if necessary, we may assume $\text{Fix}(g_*) \cap \text{Fix}((hgh^{-1})_*) = \phi$. Suppose $M \cap \partial_\infty W \neq \phi$. Then $M \cap \partial_\infty h^i(W) \neq \phi$ for every integer i as M is H -invariant. By Lemma 5.11 we have $d_*(h^i(W), \text{Fix}(g_*)) \leq 2$ for any integer i and any squared Type B isometry $g \in H$. In particular, $d_*(h^i(W), \text{Fix}(g_*)) \leq 2$ and $d_*(h^i(W), \text{Fix}((hgh^{-1})_*)) \leq 2$.

Since $d_*(\text{Fix}(h_*), \text{Fix}(g_*)) = 2$ and $d_*(W, \text{Fix}(h_*)) = 1$, Lemma 5.3 implies there are at most two integers i with $d_*(h^i(W), \text{Fix}(g_*)) = 1$. Therefore for all except at most 4 integers i we have

$$d_*(h^i(W), \text{Fix}(g_*)) = d_*(h^i(W), \text{Fix}((hgh^{-1})_*)) = 2.$$

Fix five distinct walls $W_1, W_2, W_3, W_4, W_5 \in \{h^i(W) : i = 1, 2, \dots\}$ with

$$d_*(W_i, \text{Fix}(g_*)) = d_*(W_i, \text{Fix}((hgh^{-1})_*)) = 2.$$

For each W_i ($i = 1, \dots, 5$), since $M \cap \partial_\infty W_i \neq \phi$ Lemma 5.12 implies there is a wall W'_i with

$$d_*(W_i, W'_i) = d_*(W'_i, \text{Fix}(g_*)) = d_*(W'_i, \text{Fix}((hgh^{-1})_*)) = 1.$$

Let $\tilde{W}$ be the wall such that $h_*(\tilde{W}) = \tilde{W}$ and $d_*(W, \tilde{W}) = 1$. If $W'_i = W'_j$ for some $i \neq j$, then $\tilde{W}W_iW'_iW'_j\tilde{W}$ is an injective edge loop of length 4, contradicting to Lemma 5.3. Suppose $W'_i \neq W'_j$ for $i \neq j$. Since $\text{Fix}(g_*)$ and $\text{Fix}((hgh^{-1})_*)$ are disjoint edges, $d_*(W'_i, \text{Fix}(g_*)) = d_*(W'_i, \text{Fix}((hgh^{-1})_*)) = 1$ for $i = 1, \dots, 5$ implies there are two indices $i \neq j$, and walls $W' \neq W''$ such that $g_*(W') = W'$, $(hgh^{-1})_*(W'') = W''$ and $d_*(W'_i, W') = d_*(W'_i, W'') = d_*(W'_j, W') = d_*(W'_j, W'') = 1$. Thus $W'_iW'W'_jW''W'_i$ is an injective edge loop of length 4 in G_* , a contradiction. $\square$

Lemma 5.20. *Let $M \subset \partial_\infty X$ be a H -admissible subset, $h, g \in H$ squared Type B isometries with $d_*(\text{Fix}(h_*), \text{Fix}(g_*)) = 2$, and W a wall with $d_*(W, \text{Fix}(h_*)) = 2$. Then $M \cap \partial_\infty W = \phi$.*

Proof. Since $d_*(W, \text{Fix}(h_*)) = 2$, Lemma 5.12 implies there is a set of walls $\{W_i\}_{i=1}^N$ ($N = 1$ or 2) with $d_*(W_i, \text{Fix}(h_*)) = 1$ and $M \cap \partial_\infty W \subset \cup_{i=1}^N (M \cap \partial_\infty W_i)$. Now the lemma follows from Lemma 5.19. $\square$

Lemmas 5.18, 5.19, 5.20, Proposition 5.10 and the remark before Lemma 5.13 imply that for any two squared Type B isometries $g, h \in H$, $d_*(\text{Fix}(h_*), \text{Fix}(g_*)) \neq 2$ holds. Lemma 5.17 then implies there are $h, h' \in H$ with $d_*(\text{Fix}(h_*), \text{Fix}((h')_*)) \geq 3$. Now it follows from Lemma 5.16 that H has the TA-property. The proof of Theorem 5.1 is now complete.

REFERENCES

- [AB] S. Adams, W. Ballmann, *Amenable isometry groups of Hadamard spaces*, Math. Ann. **312** (1998), no. 1, 183–195.
- [AS] U. Abresch, V. Schroeder, *Analytic manifolds of nonpositive curvature*, Collection SMF séminaires et Congrès **1** (ed. Arthur L. Besse), 1996.
- [B] W. Ballmann, *Lectures on spaces of nonpositive curvature*, volume **25** of DMV seminar. Birkhäuser, 1995.
- [BB] W. Ballmann, S. Buyalo, *Periodic rank one geodesics in Hadamard spaces*, preprint 2002.
- [BBr] W. Ballmann, M. Brin, *Orbihedra of nonpositive curvature*, Inst. Hautes Études Sci. Publ. Math. **82** (1996), 169–209.
- [BGS] W. Ballmann, M. Gromov and V. Schroeder, *Manifolds of Nonpositive Curvature*, Progress in Mathematics, Volume **61**, Birkhäuser, 1985.
- [BH] M. Bridson, A. Haefliger, *Metric spaces of nonpositive curvature*, Grundlehren **319**, Springer-Verlag, Berlin(1999).
- [Br] M. Bridson, *On the semisimplicity of polyhedral isometries*, Proceedings of the American mathematical society **127**, no. 7, 2143–2146.
- [BSw] W. Ballmann, J. Swiatkowski, *On Groups acting on Nonpositively Curved Cubical Complexes*, Enseign. Math. (2) **45** (1999), no. 1-2, 51–81.
- [CE] S. Chen, P. Eberlein, *Isometry groups of simply connected manifolds of nonpositive curvature*, Illinois J. Math. **24** (1980), no. 1, 73–103.
- [EO] P. Eberlein, B. O’Neill, *Visibility manifolds*, Pacific J. Math. **46** (1973), 45–109.
- [E1] P. Eberlein, *Euclidean de Rham factor of a lattice of nonpositive curvature*, J. Diff. Geom. **18** (1983), 209–220.
- [E2] P. Eberlein, *Lattices in spaces of nonpositive curvature*, Ann. of Math **11** (1980), 435–476.
- [E3] P. Eberlein, *Geodesic flow in certain manifolds without conjugate points*, Trans. AMS **167** (1972), 151–170.
- [G] M. Gromov, *Hyperbolic Groups*, in “Essays in Group Theory” (ed. S.Gersten) M.S.R.I. Publications No.8, Springer-Verlag(1987) 75–263.
- [H] J. Hempel, *3-Manifolds*, Ann. of Math. Studies **86**, Princeton University Press, 1976.
- [HR] C. Hruska, K. Ruane, *in preparation*.
- [HS1] C. Hummel, V. Schroeder, *Tits geometry associated with 4-dimensional closed real-analytic manifolds of nonpositive curvature*, J. Differential Geometry **48** (1998), 531–555.
- [HS2] C. Hummel, V. Schroeder, *Tits geometry of cocompact real-analytic Hadamard manifolds of dimension 4*, Differential Geometry and its Applications **11** (1999), 129–143.
- [K] M. Kapovich, *Hyperbolic Manifolds and Discrete Groups*, Progress in Mathematics, Volume **183**, Birkhäuser, 2001.
- [L] B. Leeb, *A characterization of irreducible symmetric spaces and Euclidean buildings of higher rank by their asymptotic geometry*, Bonn Mathematical Publications **326**.
- [PV] I. Pays, A. Valette, *Sous-groupes libres dans les groupes d’automorphismes d’arbres*, Enseign. Math. (2) **37** (1991), no. 1-2, 151–174.
- [R] K. Ruane, *Dynamics of the Action of a CAT(0) group on the Boundary*, Geometriae Dedicata **84** (2001), 81–99.
- [S1] V. Schroeder, *Structure of flat subspaces in low-dimensional manifolds of nonpositive curvature*, Manuscripta Math **64** (1989), 77–105.
- [S2] V. Schroeder, *Codimension one tori in manifolds of nonpositive curvature*, Geom. Dedicata **33** (1990), no. 3, 251–263.
- [Sw] E. Swenson, *A cut point theorem for CAT(0) groups*, J. Differential Geom. **53** (1999), no. 2, 327–358.
- [T] J. Tits, *Free subgroups in linear groups*, J. Algebra **20**(1972), 250–270.
- [X] Xiangdong Xie, *Groups Acting on CAT(0) Square Complexes*, preprint.

DEPARTMENT OF MATHEMATICS, WASHINGTON UNIVERSITY, ST. LOUIS, MO 63130.

E-mail address: xxie@math.wustl.edu